



NERVOUS SYSTEM *regulation*

COURSE MANUAL REVISED EDITION

NeuroSync Academy

Advanced Nervous System Regulation Course Program

Day 1

8:00 – 13:00

- *Welcoming and admin*
 - *Sign-in, CPD documentation, manuals, and housekeeping notes.*
- *Introduction*
 - *Course outline and purpose of advanced nervous system regulation.*
- *Understanding regulation*
- *Understanding the ANS – two branches*
 - *Sympathetic branch overview.*
 - *Parasympathetic branch overview.*
 - *How the two branches work together.*
- *How to assess the ANS for clinical markers*

Day 2 & 3

8:00 – 13:00

- *How to regulate the nervous system*
- *Breath*
- *Eye movements*
- *Somatic*
- *Protocol flow*
- *Three R Method: Recognize – Respond – Restore*
- *Case studies and clinical application*

(15-minute integration break each hour)



Our Vision

We believe nervous system regulation is not an “add-on,” but a foundational layer of all healing work. This course is designed to equip you with clarity, insight, confidence, and clinical precision—so that your work as a practitioner becomes more effective, more embodied, and truly transformative. By insight, we mean a deep sight from within—an inner knowing that allows you to apply this work from your own story, your own truth. Nervous system work is not something to merely understand intellectually; it must be felt first.

*Our aim is not to teach you something entirely new, but to reveal what you may already know—just in a new language, a new rhythm. So you can make it your own. Not something prescribed, but something lived.
Let’s begin.*

With appreciation,

Corinne Badenhorst

Occupational Therapist | Dipl. Psychoneuroimmunology | HeartMath Certified Anxiety & Trauma
<https://www.neurosynchronacademy.com/>



Disclaimer



@neuroSync Academy

The content presented in this course and manual is intended for educational and professional development purposes only. It is designed to support clinicians, therapists, and healthcare providers in understanding and applying principles of nervous system regulation, allostatic load, and related clinical concepts.

This material:

Does not constitute medical diagnosis, treatment, or psychological counseling. Is not intended to replace individualized medical advice, mental health evaluation, or specialist referral.

Should not be used to self-diagnose or self-treat any medical or psychiatric condition.

Presents clinical markers and symptom patterns that may suggest nervous system dysregulation, but these are intended as educational frameworks, not formal diagnostic criteria.

While many of the physiological and psychological patterns explored in this course are supported by research, they must always be interpreted within the broader context of a full clinical assessment and multidisciplinary collaboration where appropriate.

Participants are encouraged to use the tools, assessments, and frameworks taught in this course within the scope of their professional training and licensure, and to refer clients to qualified professionals when clinical complexity, safety, or diagnostic uncertainty arise.

The creators and facilitators of this course are not responsible for the application or misapplication of the material presented. This document carries copy rights. It might not be forwarded.

Introduction

A Quiet Walk Through You...

Before we begin, let's take a moment. A moment to travel back—not with judgment, but with gentleness.

Imagine you're 13 again.

It's a late afternoon. The light is golden, soft, stretching long across your bedroom wall. You've just come home from school. You're holding a folded note in your hand—the kind passed between desks, filled with squiggly hearts and scribbled hope. But today, it says something different.

He doesn't want to be your boyfriend anymore.

Your heart tightens. Your breath shortens. Something sinks. Maybe it's embarrassment. Maybe it's shame. Maybe it's that quiet voice that whispers, "I'm not enough."

You fold the note again. Smaller this time.

And you learn to fold something inside you too.

Now imagine the years roll forward.

One moment here. Another there.

An adult raises their voice. A friend disappears.

You learn to hold your breath before speaking.

You study harder to prove you're worth noticing.

You don't say what hurts—you smile instead.

Bit by bit, your body adapts.

Your nervous system, brilliant and loyal, does what it must:
tightens, sharpens, guards, performs.

This is the invisible weight we carry.

Not just the big traumas—but the everyday moments that taught your system how safe it is—or isn't—to be fully you.

This weight has a name: allostatic load.

The cost of constantly adjusting to stress.

Over time, it shifts your baseline.

You get used to running on edge.

You forget what calm feels like.

You start making choices—not from presence, but protection.

And this... this is why nervous system regulation matters.

Not just for our clients.

But for us.

So we can remember what it feels like to live from truth, not tension.

Before we go into the science, the maps, and the techniques—
just take a moment here.

Breathe.

Notice what your 13-year-old self might still be holding.

Not to fix it.

Just to see it.

Because healing doesn't begin with doing.

It begins with noticing.

And this is why this course exists.

Because beneath every technique, every client session, and every therapeutic approach—there is a nervous system trying to find safety.

This course was born from a deep need to bridge insight with embodiment...

To go beyond understanding trauma & dysregulation and into transforming it.

It exists because you, as a practitioner, deserve to work from a place that's not just informed—but integrated.

And because your clients don't just need support.

They need you—your presence, your nervous system, your steady ground.

Every day, clients walk into your practice carrying untold stress stories—woven into their tissues, echoed in their breath, posture, digestion, and emotional patterns. Too often, talking alone can't reach the layers where those stories live.

We all carry stories. And like the Hebrew language, they often only make sense when read in reverse—when we look back with insight, presence, and compassion.

This course is here to help you do just that.

It was created for therapists, clinicians, and wellness practitioners who want more than insight—they want embodied clinical skill. Tools that help clients regulate from the inside out.

That means knowing the language of the body, the brain, and the breath—and being able to speak it fluently, with precision and care.

This isn't a course of quick fixes or superficial techniques. It's about learning the truth beneath the "how-tos." It's about transformation—from the roots.

Imagine someone had told you at age 13 that there was a safe space inside you where your story lived—and that this space could become a place of healing, not just survival. What might have changed?

Survival vs. Creation: Two States, Two Realities

When we live in survival states—fight, flight, freeze, or fawn—the body contracts, the mind loops, and the soul goes quiet. We react, adapt, and perform. We live from the outside in.

But when we shift into states of creation and flow, the nervous system opens. Breath deepens. Thoughts integrate. We become responsive, not reactive. The body softens, the brain synchronizes, and the soul begins to speak again.

This course is about helping you—and your clients—move from a reactive life to a consciously lived one.

The End Goal: We You'll Be Able to Do

By the end of this course, you'll be able to:

- Accurately identify signs of nervous system dysregulation
- Understand root causes—biological, emotional, and somatic
- Choose and apply the correct regulation techniques
- Sequence interventions for maximum therapeutic impact
- Track client shifts and adapt protocols in real time

Why This Workbook Matters

This isn't just a note-taking companion.

It's a clinical thinking tool.

A place to process, practice, reflect, and embody the knowledge you're gaining—so that you don't just learn nervous system regulation...

You *live* it.

The Why behind the why...

Nervous System Regulation Is a possible missing Link in Modern Healthcare

I didn't set out to become an expert in nervous system regulation.

I was pulled into it—by trauma, by loss, by moments when no technique, no therapy, and no diagnosis could explain what was happening inside me.

My nervous system was speaking long before I knew its language.

It was through my own unraveling that I began to see the missing link—not just in myself, but in clients, in colleagues, in the broader world of healthcare.

And then the science caught up with what I already knew in my bones.

1. Before cognition, before emotion, before behavior — there is state.

The autonomic nervous system (ANS) determines whether your client can:

- Think clearly
- Form healthy attachments
- Digest food and absorb nutrients
- Regulate hormones and sleep
- Or spiral into anxiety, inflammation, IBS, or autoimmune flare

Therapy, medication, or lifestyle advice can't stick if the nervous system is saying "I'm not safe."

2. Regulation restores physiological coherence.

When HRV improves, vagal tone rises, and the system finds rhythm again, we don't just feel better—we function better:

- Immune response strengthens¹
- Pain thresholds shift²
- Sleep quality improves³
- Cognitive flexibility and prefrontal control return⁴
- Even blood sugar regulation stabilizes⁵

This is the biology of safety.

3. Dysregulation can mimic psychiatric illness.

Many clients labeled with:

- "Treatment-resistant depression"
- "Borderline personality disorder"
- "Panic disorder"

...are often stuck in looping dysregulated autonomic states—not resistant, not disordered, just trying to survive.

4. Regulation isn't just symptom management. It's root restoration.

You're not just giving people tools to "calm down."

You're restoring their body's capacity to:

- Process experience
- Self-regulate
- Connect
- Recover
- Live

It's not soft. It's clinical neuroscience.

It's not fringe. It's the foundation.

This course stands at the intersection of personal truth and scientific integrity—because both are necessary for real healing.

In short: Nervous system regulation is not an "add-on" to therapy or medicine. It's the **platform** every other intervention stands on. Without it, we're just rearranging the furniture in a burning house.

References

1. Tracey KJ. The inflammatory reflex. *Nature*. 2002;420(6917):853-859.
2. Payne P, Levine PA, Crane-Godreau MA. Somatic experiencing: using interoception and proprioception as core elements of trauma therapy. *Front Psychol*. 2015;6:93.
3. Porges SW. *The Polyvagal Theory: Neurophysiological Foundations of Emotions, Attachment, Communication, Self-Regulation*. New York: Norton; 2011.
4. Thayer JF, Åhs F, Fredrikson M, Sollers JJ 3rd, Wager TD. A meta-analysis of heart rate variability and neuroimaging studies: implications for heart rate variability as a marker of stress and health. *Neurosci Biobehav Rev*. 2012;36(2):747-756.
5. Carnevali L, Sgoifo A. Vagal modulation of resting heart rate in rats: the role of stress, psychosocial factors, and physical exercise. *Front Physiol*. 2014;5:118.

Before Insight Comes Regulation

Most therapeutic models begin with cognition—thought tracking, meaning-making, insight-building.

But here's the truth:

If the nervous system is dysregulated, no amount of insight will land.

Insight cannot integrate if the body still feels unsafe.

Why Regulation Must Come First

Bottom-Up Before Top-Down

Trauma, anxiety, burnout, and chronic illness don't start in faulty thoughts—they start in body-brain loops.

When the autonomic nervous system is activated, the prefrontal cortex (PFC)—your center for reasoning and reflection—goes offline.

That's why therapeutic work must begin with tools that speak the body's language:

Breath. Rhythm. Posture. Tone. Movement.

We work bottom-up before we work top-down.

Rhythm Over Reason

The nervous system doesn't speak logic.

It speaks pattern and pace—like a musical instrument.

Regulation isn't just about calming down.

It's about restoring rhythmic flexibility—the ability to shift gears, move between states, and return to center.

The goal is not stillness.

The goal is flow and Coherence is the Gateway...

Healing begins when the system enters coherence—

a state where breath, heart, brain, and vagus nerve move in sync.

This is the threshold where safety becomes biological.

Where the body says, "I can begin again."

That's why we don't start with insight.

We start with regulation.

Because before a client can heal, they must first feel safe enough to stay.

*"You
cannot
talk a
survival
system into
safety.
You must
regulate
it."*

Why Regulation First— Quick Summary

A clinical summary of what changes when safety comes first.

Area of Function	Without Regulation	With Regulation
Cognitive Function	Foggy thinking, rumination, PFC shutdown	Clear thinking, increased focus, cognitive flexibility
Emotional Processing	Flooding, shutdown, looping reactions	Emotional range, responsiveness, integration
Digestive & Hormonal Health	Bloating, IBS, adrenal dysregulation, hormonal swings	Improved gut motility, cortisol rhythm, endocrine balance
Sleep & Recovery	Insomnia, night waking, light sleep	Deeper rest, parasympathetic dominance at night
Immune Function	Chronic inflammation, poor healing, flare-ups	Stronger immune modulation, anti-inflammatory vagal tone
Attachment & Relationships	Fawn, disconnect, hypervigilance, mistrust	Safety in connection, healthy boundaries, attuned relating
Therapy & Learning	Insight doesn't stick, resistance, overwhelm	Integration of insight, capacity to change, neuroplasticity
Long-term Health	High allostatic load, chronic disease risk	Systemic resilience, homeostasis, adaptability

Without safety, the system protects.

With safety, the system transforms.

References

- Tracey KJ. The inflammatory reflex. *Nature*. 2002;420(6917):853-859.
- Payne P, Levine PA, Crane-Godreau MA. Somatic experiencing: using interoception and proprioception as core elements of trauma therapy. *Front Psychol*. 2015;6:93.
- Porges SW. *The Polyvagal Theory: Neurophysiological Foundations of Emotions, Attachment, Communication, Self-Regulation*. New York: Norton; 2011.
- Thayer JF, Åhs F, Fredrikson M, Sollers JJ 3rd, Wager TD. A meta-analysis of heart rate variability and neuroimaging studies: implications for heart rate variability as a marker of stress and health. *Neurosci Biobehav Rev*. 2012;36(2):747-756.
- Carnevali L, Sgoifo A. Vagal modulation of resting heart rate in rats: the role of stress, psychosocial factors, and physical exercise. *Front Physiol*. 2014;5:118.

What This Course Will Teach You

This course equips therapists, clinicians, and wellness professionals to:

- **Work with the body first**

Learn to read and respond to physiological state before cognition or conversation.

Because story doesn't integrate until safety is felt.

- **Use rhythm as a clinical tool**

Harness the power of breath, pacing, sound, and movement to regulate the nervous system and restore flexibility.

- **Build protocols that stabilize**

Create personalized sequences that anchor clients in safety before deep work begins—preventing overwhelm, retraumatization, or shutdown.

- **Prioritize state before exploring story**

Learn to map your client's nervous system state in real time, and shift it—so that when the story is told, the body is ready to receive it.

The Quiet Overflow

Every day, there are drips.

Deadlines. Missed meals. Family drama.

Financial tension. Traffic. That difficult co-worker. The thing you forgot to say.

None of it seems like a big deal on its own.

But what if we keep adding,
without ever letting anything out?

We dysregulate.

We burn out.

We shut down.

We panic.

Not always because of one big trauma—
but because we never emptied.

We never returned to baseline.

And baseline is where it all waits:

Your identity.

Your purpose.

Your immune system.

Your dreams.

Your story.

This is what regulation makes possible.

And this is what you'll learn to offer others.

*"Safety is the treatment.
Regulation is the access
point.
Rhythm is the medicine."*

Caught in the Cage: A Reflection

*Imagine being trapped in a cycle of fear.
Not a dramatic panic—but a quiet tightening.
A hum beneath the surface. A subtle narrowing of breath.
It starts slow. You over-function. You please. You avoid.
You do what's expected. You lose your appetite, or your voice, or your joy.
And the world shrinks.
Not because it is small—but because your system is on alert.
Your perception shifts from freedom to survival.
This isn't just emotion.
It's energy in motion.
It's your nervous system adapting to what it believes is threat.
And then comes the effort...
A thousand ways of managing symptoms.
A thousand triggers and traps to avoid.
As poet Gillian Gibbons writes of fear:
"Fear is the cheapest room in the house. I would like to see you in better living conditions."
This course is that invitation.
Not just to learn—
but to regulate, to return, to reclaim your spaciousness.
Because as Rumi once said:
"Stop acting so small. You are the universe in ecstatic motion."
So before we go deeper—
stop.
Listen.
Breathe.
Experience what it means to come home to your baseline.*

Experience Your Own Baseline

Before we move deeper into the course, let's pause—not to think, but to feel.

You've just read about baseline.

How our system accumulates without release.

How the body can't heal if it never returns to stillness.

Now, let's give your nervous system a chance to feel...

<https://music.apple.com/za/album/mamuna-papua/1763305994?i=1763305995>

1 Before Listening — “Baseline Check-In”

Current physical state:

- My breath feels:
 - ☐ Fast ☐ Slow ☐ Tight ☐ Open ☐ Don't know
- My heart feels:
 - ☐ Heavy ☐ Calm ☐ Fast ☐ Tight
- My muscles feel:
 - ☐ Tense ☐ Relaxed ☐ Jumpy ☐ Numb
- My thoughts are:
 - ☐ Racing ☐ Still ☐ Foggy ☐ Focused

2 While Listening — “The Body’s Conversation”

Where do you sense the music most?

- ☐ Head
- ☐ Chest / Heart
- ☐ Belly / Gut
- ☐ Hands
- ☐ Legs
- ☐ Other: _____

What sensations arise?

- ☐ Warmth
- ☐ Tingling
- ☐ Heaviness
- ☐ Floating
- ☐ Movement urge (swaying, tapping, breath changes)
- ☐ Release (yawning, sighing, tears)

Emotional tone (non-verbal):

- ☐ Safe
- ☐ Open
- ☐ Soft
- ☐ Connected
- ☐ Alert
- ☐ Grounded
- ☐ Grateful
- ☐ Tender

3 After Listening — “Integration Reflection”

What changed?

- ☐ My breathing is easier
- ☐ My muscles softened
- ☐ My thoughts slowed
- ☐ My emotions feel more accessible
- ☐ I feel more in my body

You Can't Teach What You Haven't Felt

*This moment of music is more than a pause—
it's a personal entry point into the work we'll be
doing with others.*

*You've just met your own nervous system in
rhythm.*

Let's continue from here.

*If you could name your nervous system state
now in one word, what would it be?*

Nervous System Insight

As you listened, your brainstem, vagus nerve,
and limbic system received input far faster than
your thinking brain.

Regulation starts before cognition...

Part 1

Foundations of Nervous System Regulation

"We treat the state, not just the symptom."

The Antidote Is Rhythm

"Regulation is not about doing less. It's about returning to rhythm—like the breath, like the tide, like the heart that knows its beat."

What We're Really Trying To Do

We're not trying to avoid stress, avoid emotion, or "stay calm."

We're trying to restore rhythm.

We're trying to bring the system back into flow.

Dysregulation happens when the orchestra of the nervous system loses its timing.

The players—the heart, brain, gut, hormones, and breath—start moving to different rhythms.

Balance is not silence. It is synchrony.

And that's what regulation is:

A return to physiological coherence.

Where signals between brain, heart, and body start harmonizing again.

Nervous system regulation refers to the body's ability to *move fluidly* between various autonomic states *in response to context, and to return to a baseline of safety* and rest when the stressor is removed. This adaptability is critical for physical health, psychological stability, and social engagement.

It is a dynamic process involving brain-body communication via the autonomic nervous system (ANS), which governs not only obvious survival responses (e.g. fight, flight, freeze), but also subtle physiological functions such as:

- Digestion
- Heart rate variability (HRV)
- Pupil dilation
- Hormone release
- Emotional tone

When Well-Regulated

- Clear thinking and focus
- Appropriate emotional expression
- Stable digestion, sleep, and hormone cycles
- Resilient stress recovery
- Connection and curiosity in social interactions

When Dysregulated

Individuals may experience symptoms such as:

- Anxiety
- Irritability
- Fatigue
- Chronic pain
- Digestive disorders
- Emotional numbness

References

1. Elbers J, Batista M. Treating nervous system dysregulation in children. *Cdn J Cont Med Ed*. 2018.
2. Thayer JF, Lane RD. Claude Bernard and the heart-brain connection: further elaboration of a model of neurovisceral integration. *Neurosci Biobehav Rev*. 2009;33(2):81-88.
3. Perry BD, Szalavitz M. *The Boy Who Was Raised as a Dog: And Other Stories from a Child Psychiatrist's Notebook*. New York: Basic Books; 2017.
4. Porges SW. *The Polyvagal Theory: Neurophysiological Foundations of Emotions, Attachment, Communication, Self-Regulation*. New York: Norton; 2011.
5. McCraty R, Zayas MA. Cardiac coherence, self-regulation, autonomic stability, and psychosocial well-being. *Front Psychol*. 2014;5:1090.
6. Payne P, Levine PA, Crane-Godreau MA. Somatic experiencing: using interoception and proprioception as core elements of trauma therapy. *Front Psychol*. 2015;6:93.

The Physiology Behind Regulation

The Autonomic Nervous System (ANS): The Body's Regulatory Core

The body doesn't just respond to the mind—it informs it.

A Brief History: What We Missed

In the early 1900s, physiologist J.N. Langley discovered the autonomic nervous system (ANS)—the system that controls breath, heart rate, digestion, and the stress response.

It was a revolutionary insight.

But it was only half the story.

For decades, the dominant model assumed the brain was the master, and the body the servant.

So mental health interventions focused on cognition, medication, and behavior—what we could see and think.

But regulation doesn't begin in the mind.

It begins in the state beneath thought.

It begins in the body.

As trauma researcher Dr. Bessel van der Kolk later wrote:

"The body keeps the score."

Regulation Starts with the ANS

The autonomic nervous system is our body's internal regulation system.

It tells us whether it's safe to connect—or time to protect.

It's not controlled by conscious thought.

It's an ancient, adaptive intelligence that keeps us alive.

It has two main branches:

1. Sympathetic Nervous System (SNS) The Activator

- Mobilizes the body under stress: fight or flight
- Increases heart rate and blood pressure
- Heightens vigilance and muscular tension
- Suppresses digestion, reproduction, and long-term repair
- Essential in short-term danger—harmful when stuck "on"

2. Parasympathetic Nervous System (PNS) The Restorer

- Calms the body and restores homeostasis
- Slows heart rate, deepens breath
- Activates digestion, immune repair, hormonal rhythm
- Centered in the vagus nerve—especially the ventral vagal branch, which supports connection and social engagement
- Healing happens here. Regulation lives here.

Why This Matters in Practice

When we understand the ANS, we stop chasing symptoms.

We begin to listen to the state underneath the story.

Dysregulation is not a disorder.

It is a signal that the system doesn't feel safe.

Regulation snapshot

State	Physiological Markers	Clinical Implications
Regulated	Stable HRV, rhythmic breath, social cues active	Calm alertness, digestion, engagement
Sympathetic	Rapid HR, shallow breath, tense posture	Anxiety, irritability, overdrive
Dorsal Vagal	Low HRV, slowed digestion, flat affect	Numbness, depression, withdrawal

1. Langley JN. The autonomic nervous system. Brain. 1903;26(1):1-26.

2. Porges SW. The polyvagal perspective. Biol Psychol. 2007;74(2):116-43.

3. van der Kolk BA. The Body Keeps the Score. New York: Penguin; 2015.

4. Thayer JF, Lane RD. Claude Bernard and the heart-brain connection. Neurosci Biobehav Rev. 2009;33(2):81-88.

5. McEwen BS. Physiology and neurobiology of stress and adaptation. Physiol Rev. 2007;87(3):873-904.

6. Payne P, Levine PA, Crane-Godreau MA. Somatic experiencing: using interoception and proprioception as trauma tools. Front Psychol. 2015;6:93.

Take a micro-pause and see if you could recognise your
own nervous system state right now...
Close your eyes for a moment and focus on the sensations in your
body and your breath.

Part 2

Understanding the bigger picture

Everything we will explore in this course lives inside one simple truth:

You cannot heal what you cannot feel.

You cannot rewire what you cannot first regulate.

Think of nervous system work like a tree

The Roots: RECOGNIZE

Awareness is the soil.

Before change, we must see and name what state the nervous system is in — safety, survival, or shutdown.

This is where we pause, notice, and listen.

The Trunk: RESPOND

Response is the lifeline.

Once we recognize the state, we meet it with the right tools—
breath, movement, rhythm, and tone.

Here the system begins to shift, and flexibility returns.

The Branches: RESTORE

When the system feels safe and regulated, deeper healing begins.

Patterns can change.

Stories can integrate.


The body, brain, and soul can return to wholeness.

Every exercise, every protocol, every reflection in this course is simply helping you (and your clients) climb this tree — from roots, through the trunk, into the branches.

Without roots, the tree cannot grow.

Without the trunk, it cannot stand.

Without branches, it cannot reach.



Meet the Autonomic Nervous System

“The body’s hidden conductor of protection, connection, and collapse.”

What Is the ANS?

The autonomic nervous system (ANS) is the part of the nervous system that controls involuntary bodily functions like heart rate, digestion, breathing, hormone release, immune function, and even pupil dilation.

It operates below conscious awareness and determines whether we are in a state of safety, mobilization, or shutdown.

The state of your nervous system dictates how you feel, how you relate, how you learn, how you heal — and whether your body feels like home or like a battlefield.

The ANS Has Two Main Branches

Healthy regulation means the ANS shifts fluidly between these branches depending on context and perceived safety.

- Sympathetic Nervous System (SNS)
- → Activates the body for action.
- Think: stress, alertness, mobilization (fight/flight).
- Parasympathetic Nervous System (PNS)
- → Calms the system and restores balance.
- Think: rest, digestion, repair, connection.

The Parasympathetic System Has Two Pathways

(Polyvagal Theory – Dr. Stephen Porges)

The PNS isn't a single pathway — it has two distinct branches of the vagus nerve that determine how we respond to safety or threat.

Understanding How the ANS Responds

The ANS doesn't respond to logic, reasoning, or words—it responds to patterns of safety and threat detected by the body. This unconscious system, called neuroception (Porges, 2004), constantly scans your internal and external environment to decide whether to mobilize, shut down, or regulate.

Before we explore deeper brain structures, it's important to understand how this survival-first wiring shapes therapy outcomes.

Clinical Insight:

“The ANS does not ask, ‘Is this true?’
It asks, ‘Am I safe?’”

Understanding which branch is currently dominant gives you a diagnostic compass for matching therapy to biology.

Key Brain Regions That Drive ANS Responses

These areas work together to prioritize survival first, and only later allow healing, learning, or connection.

Brain Region	Role in ANS Regulation
Brainstem	Receives sensory input and directs autonomic output
Hypothalamus	Coordinates endocrine, immune, and autonomic response
Insula & ACC	Interoception and emotion regulation
Amygdala	Threat detection and fear response

References

1. Porges SW. The polyvagal perspective. *Biol Psychol.* 2007;74(2):116-143.
2. Thayer JF, Lane RD. A model of neurovisceral integration in emotion regulation. *Neurosci Biobehav Rev.* 2000;24(3):225-229.
3. Taylor KS, Seminowicz DA, Davis KD. Two systems of resting state connectivity between the insula and cingulate cortex. *Hum Brain Mapp.* 2009;30(9):2731-2745.
4. Critchley HD, Harrison NA. Visceral influences on brain and behavior. *Neuron.* 2013;77(4):624-638.
5. Smith R, Lane RD, Nadel L, Moutoussis M. A computational neuroscience perspective on the change process in psychotherapy. *Neuropsychol Rev.* 2020;30(3):274-297.

The Sympathetic Nervous System:

Anatomy of Hyperarousal

Anatomy of Hyperarousal – When Survival Gets Stuck On

What Is the Sympathetic Nervous System?

The sympathetic nervous system (SNS) is the branch of the ANS responsible for activating the body's fight-or-flight response.

It's designed to help you respond to danger — but when chronically activated, it becomes the biology of burnout, anxiety, and inflammation.

Where It Starts: Thoracolumbar Activation

Unlike the vagus nerve (which originates in the brainstem), the SNS begins in the thoracolumbar region of the spinal cord — segments T1 to L2.

Here's how the activation flows:

1. Threat is detected by the amygdala
2. The hypothalamus sends a signal
3. Sympathetic neurons fire from the spinal cord
4. Messages travel via the sympathetic chain (paravertebral ganglia)
5. Target organs receive the message:
6. Mobilize. Survive. Now.

Neurotransmitters and Hormones

At each target site, the SNS releases:

- Norepinephrine (noradrenaline): sharpens focus, increases heart rate
- Epinephrine (adrenaline): released from adrenal medulla into bloodstream
- Cortisol: triggered via the HPA axis for prolonged stress response

Short-term, these chemicals protect us.

Long-term, they can wear down immunity, alter hormones, trigger inflammation, and flood the brain with anxiety loops.

The Sympathetic Nervous System: Clinical markers

When the sympathetic branch of the autonomic nervous system (ANS) dominates, the body mobilizes for survival. This “fight-flight” state is adaptive short-term, but when prolonged it creates chronic dysregulation, impacting physical, emotional, and behavioral health.

Physiological Markers

- ↑ Heart rate / palpitations – resting tachycardia or sudden spikes.
- ↑ Respiratory rate – shallow, upper-chest breathing, frequent sighs.
- ↑ Blood pressure – sustained elevation.
- Dilated pupils – mydriasis for increased scanning.
- Sweating – clammy palms, underarm sweating, damp hairline.
- Cold hands and feet – peripheral vasoconstriction.
- Skin color shifts – flushed cheeks, blotchy neck/chest, or sudden paleness (due to blood redistribution).
- Goosebumps (piloerection) – often on arms or back of neck.
- Dry mouth – reduced saliva flow.
- GI suppression – nausea, “butterflies,” constipation, or urge to empty bowels.

Facial & Micro-Markers

- Jaw tightness – clenched jaw, teeth grinding (bruxism).
- Lips – pressed together tightly or thinned out; corners turned down.
- Nostrils – slightly flared during alert scanning.
- Eyebrows – drawn together, forming vertical “worry lines.”
- Eye movements – scanning the room, darting side to side.
- Facial pallor or redness – alternating flush/pale states depending on adrenaline surges.

Posture & Motor Markers

- Rigid, high-tension posture – shoulders raised, chest lifted.
- Fidgeting / restlessness – tapping fingers, bouncing leg, pacing.
- “Ready-to-spring” stance – weight slightly forward, knees flexed.
- Tremors or micro-shakes – especially in hands.

Voice & Communication Markers

- Voice pitch – higher, thinner tone when speaking.
- Speech rate – talking too fast or clipped, abrupt sentences.
- Vocal tension – cracking, straining, or “tight” sound.
- Interruptions – frequent sighing, swallowing, or throat clearing.

Cognitive & Emotional Markers



- Racing thoughts, mental rehearsal of danger.
- Black-and-white thinking, impulsive decisions.
- Anxiety, irritability, frustration, or sudden anger.
- Hypervigilance and exaggerated startle response.

Behavioral Markers

- Compulsive scanning/checking (doors, phones).
- Over-preparing, over-functioning (“doing” to avoid “feeling”).
- Startle response to small noises or movements.
- Sleep disruption – difficulty falling asleep or waking wired.
- Food changes – loss of appetite or stress eating.

References

- Thayer JF, Åhs F, Fredrikson M, Sollers JJ, Wager TD. A meta-analysis of heart rate variability and neuroimaging studies: Implications for heart rate variability as a marker of stress and health. *Neurosci Biobehav Rev.* 2012;36(2):747–756.
- Jerath R, Edry JW, Barnes VA, Jerath V. Physiology of long pranayamic breathing: Neural respiratory elements may provide a mechanism that explains how slow deep breathing shifts the autonomic nervous system. *Med Hypotheses.* 2006;67(3):566–571.
- Esler M, Lambert E, Schlaich M. Chronic activation of the sympathetic nervous system is the dominant contributor to systemic hypertension. *J Appl Physiol.* 2010;109(6):1996–1998.
- McCorry LK. Physiology of the autonomic nervous system. *Am J Pharm Educ.* 2007;71(4):78.
- Critchley HD. Neural mechanisms of autonomic, affective, and cognitive integration. *J Comp Neurol.* 2002;493(1):154–166.
- Arnsten AF. Stress signalling pathways that impair prefrontal cortex structure and function. *Nat Rev Neurosci.* 2009;10(6):410–422.
- van der Kolk B. *The Body Keeps the Score.* New York: Viking; 2014.
- Pole N. The psychophysiology of posttraumatic stress disorder: A meta-analysis. *Psychol Bull.* 2007;133(5):725–746.
- Anderson B, Blankenbaker D, Hanrahan S. Stress, muscle tension, and pain. *Clin J Pain.* 2014;30(10):855–862.
- Mayer EA. The neurobiology of stress and gastrointestinal disease. *Gut.* 2011;60(2):307–317.



Sympathetic Innervation

Now that we understand the body's arousal system, let's meet its equal and opposite partner – the system of calm, repair, and connection.

Parasympathetic Nervous System: The Vagus Nerve

Gateway to Safety and Shutdown

“One nerve. Two pathways. Infinite survival responses.”

What Is the Vagus Nerve?

The vagus nerve is the 10th cranial nerve and the primary nerve of the parasympathetic nervous system.

It connects the brainstem to nearly every major organ: heart, lungs, digestive tract, liver, kidneys, vocal cords, and more.

Its name comes from the Latin *vagus*, meaning “wandering”—and indeed, it wanders throughout the body, collecting information and regulating key functions.

The Vagus Regulates:

Heart rate and blood pressure

Breath rhythm

Digestion and peristalsis

Inflammation and immune response

Facial expression, voice tone, and eye contact

Emotional regulation and trauma recovery

More than 80% of its fibers are afferent, meaning they send messages from the body to the brain—not the other way around. The body speaks first. The brain responds second.

When the vagus nerve is functioning well, clients can connect, recover, and integrate.

Origin of the Parasympathetic Nervous System (PNS)

The parasympathetic branch of the autonomic nervous system arises from two main regions:

The craniosacral outflow:

- Cranial origin: Brainstem nuclei of cranial nerves — primarily the vagus nerve (cranial nerve X), but also III (oculomotor), VII (facial), and IX (glossopharyngeal).
- Sacral origin: Spinal cord segments S2-S4, responsible for pelvic organs and lower digestion.

Together, this system regulates the body's rest, digest, restore, and social connection functions.

Vagus Nerve – The Primary Parasympathetic Pathway

The vagus nerve (CN X) is the main parasympathetic highway, accounting for approximately 75–80% of parasympathetic output. It exits the medulla in the brainstem and travels downward, innervating the heart, lungs, throat, vocal cords, esophagus, stomach, liver, pancreas, intestines, and more.

The vagus is bidirectional — with 80% of its fibers carrying information from body to brain (afferent), and only 20% descending from brain to body (efferent).

It is subdivided into two key branches (per polyvagal theory):


- Ventral vagal complex: Originates in the nucleus ambiguus, supports calm, connection, facial expressivity, and vocal prosody.
- Dorsal vagal complex: Originates in the dorsal motor nucleus of the vagus, slows metabolism and heart rate, and activates freeze or shutdown when overstimulated.

Chemical Messengers



Unlike the sympathetic system (which uses norepinephrine and epinephrine), the parasympathetic system primarily uses acetylcholine (ACh) — a neurotransmitter that slows the heart, stimulates digestion, and promotes relaxation and repair.

The vagus isn't just about rest.

It's about discernment — knowing when to connect, when to slow down, and when to step back.



Parasympathetic Innervation



Two Vagal Pathways, Two Survival Outcomes (Polyvagal Theory)

According to Dr. Stephen Porges’ Polyvagal Theory, the parasympathetic system is not a single pathway, but two distinct branches of the vagus nerve

The parasympathetic nervous system is not a single road.

The vagus nerve divides into two distinct branches, each with its own origin, purpose, and survival response:

Branch	Origin	Function	State
Ventral Vagal Complex	Brainstem (nucleus ambiguus)	Social engagement, calm, connection	Safe & connected
Dorsal Vagal Complex	Lower brainstem (dorsal motor nucleus)	Energy conservation, shutdown, dissociation	Collapse, freeze, numb

The vagus nerve has two distinct branches, each with its own origin, role, and state.

The ventral vagal complex originates in the nucleus ambiguus in the brainstem. This branch supports calm, connection, and co-regulation. When the ventral vagal system is active, clients breathe more slowly and evenly, their faces and voices become expressive, and they are able to make eye contact. They feel grounded, present, and able to connect with others. *Clinically, this is the state where therapy works best – insight lands, emotional processing feels safe, and healing can begin to integrate.*

The dorsal vagal complex originates in the dorsal motor nucleus of the vagus. This branch is designed to conserve energy, slow metabolism, and activate a freeze or shutdown response when the system feels overwhelmed. When dorsal vagal dominance takes over, clients may appear blank or distant, their posture heavy, their voice flat, and their facial expression absent. They might faint, withdraw, or describe feeling hopeless, empty, or like they “aren’t really here.” *Clinically, this is the physiology of shutdown. Pushing a client for deep processing in this state risks retraumatizing them. Instead, they need slow, gentle interventions – sensory grounding, breath pacing, and co-regulation – to bring them back into a state of safety.*

References

1. Porges SW. The polyvagal theory: Neurophysiological foundations of emotions, attachment, communication, and self-regulation. Norton; 2011.
2. Dana D. The Polyvagal Theory in Therapy. Norton; 2018.
3. Kline R. Clinical applications of the vagus nerve in trauma therapy. J Mind-Body Integration. 2020;4(1):15-22.
4. Dr Andrew Huberman – youtube video

The Parasympathetic Nervous System:

Clinical markers

(Ventral Vagal – Rest, Repair, Restore)

The ventral vagal branch of the parasympathetic nervous system supports social engagement, regulation, and healing. When active, it creates a state of safety and connection that allows the body to rest, repair, and restore.

Physiological Markers

- Heart rate – moderate, stable, within healthy baseline range (Thayer et al., 2012).
- Respiration – slow, diaphragmatic, smooth; nasal breathing common (Jerath et al., 2006).
- Blood pressure – stable and often slightly lowered (McCorry, 2007).
- Digestive activity – audible bowel sounds (“rest-and-digest”), normal appetite, improved GI motility (Mayer, 2011).
- Skin tone – warm, even color; cheeks slightly pink, no blotching (Porges, 2011).
- Muscle tone – loose, flexible, not collapsed.
- Salivation – moist mouth, easier swallowing.
- Immune support – parasympathetic state enhances vagal anti-inflammatory pathways (Tracey, 2002).

Facial & Micro-Markers

- Eyes – soft gaze, eyes not darting; full “eye contact without strain.”
- Eyebrows – relaxed, not furrowed.
- Jaw & lips – neutral or slight upward curve; jaw loose, no grinding.
- Forehead – smooth (no frown lines).
- Skin – natural glow; warmth visible in face and hands.

Posture & Motor Markers

- Open posture – shoulders gently back and down, chest open.
- Movements – fluid, unforced, natural gestures.
- Stillness – patient can rest without jitteriness or collapse.

Voice & Communication Markers

- Voice tone – warm, resonant, melodic (“prosody”) (Porges, 2011).
- Speech – steady pace, natural rhythm, full sentences.
- Laughter / lightness – spontaneous smiles and humor surface easily.

Cognitive & Emotional Markers

- Calm, present-focused awareness.
- Creativity, curiosity, and perspective-taking.
- Emotional stability – can experience joy, gratitude, compassion.
- Improved focus, problem-solving, and decision-making.

Behavioral Markers

- Natural pauses and rest without guilt or collapse.
- Comfort with connection – eye contact, gentle touch, conversation.
- Engagement in social interaction – able to play, joke, and share.
- Healthy appetite and steady sleep patterns (restorative sleep, not “shut-down” sleep).

Clinical Note for Therapists

When the ventral vagal system is engaged, the client’s body “feels safe.” You’ll notice softening: the jaw unclenches, voice warms, gestures loosen. They don’t just stop bracing – they start engaging. This state is where therapy sticks: memory reconsolidates, neuroplasticity opens, and healing accelerates.

References

1. Thayer JF, Åhs F, Fredrikson M, Sollers JJ, Wager TD. A meta-analysis of heart rate variability and neuroimaging studies: Implications for heart rate variability as a marker of stress and health. *Neurosci Biobehav Rev.* 2012;36(2):747-756.
2. Jerath R, Edry JW, Barnes VA, Jerath V. Physiology of long pranayamic breathing: Neural respiratory elements may provide a mechanism that explains how slow deep breathing shifts the autonomic nervous system. *Med Hypotheses.* 2006;67(3):566-571.
3. McCorry LK. Physiology of the autonomic nervous system. *Am J Pharm Educ.* 2007;71(4):78.
4. Mayer EA. The neurobiology of stress and gastrointestinal disease. *Gut.* 2011;60(2):307-317.
5. Porges SW. *The Polyvagal Theory: Neurophysiological Foundations of Emotions, Attachment, Communication, and Self-Regulation.* New York: Norton; 2011.
6. Tracey KJ. The inflammatory reflex. *Nature.* 2002;420:853-859.

The Parasympathetic Nervous System:

Clinical markers

(Freeze • Collapse • Immobilization)

The dorsal vagal branch of the parasympathetic system is the “emergency brake”: it conserves energy by shutting the system down when threat feels inescapable. Short bursts can protect in extreme danger — but chronic dorsal activation leads to collapse, dissociation, and loss of vitality.

Physiological Markers

- Low heart rate (bradycardia) – can be deceptively “calm,” but reflects shutdown (Thayer et al., 2012).
- Shallow or slowed breathing – sometimes breath-holding; sighing or gasping.
- Low blood pressure – dizziness, fainting spells (Levine, 2015).
- Pale/ashen skin – reduced blood flow to surface; “greyed-out” complexion.
- Cold skin & extremities – strong vasoconstriction.
- Slack muscle tone – heavy limbs, “melting” posture (Mayer, 2011).
- GI changes – nausea, bloating, slowed motility.
- Low voice volume – faint, monotone voice.
- Energy collapse – fatigue, feeling “drained” or immobilized.

Facial & Micro-Markers

- Flat facial affect – little expression; blank or distant stare.
- Eyes – glassy, unfocused, or downcast; slow blinking or “frozen” gaze.
- Jaw & lips – slack mouth, corners downturned; sometimes slightly open.
- Skin – may appear greyish, bluish, or “washed out.”

Posture & Motor Markers

- Slumped posture – head down, shoulders rounded forward.
- Minimal movement – heaviness, frozen stillness.
- Collapse responses – fainting, buckling knees, or sliding down chair.

Voice & Communication Markers

- Voice tone – flat, monotone, quiet.
- Speech – slow, long pauses, reduced prosody.
- Words – minimal responses (“I don’t know,” “It doesn’t matter”).

Cognitive & Emotional Markers

- Numbness – “I can’t feel anything.”
- Dissociation – detachment from body or surroundings.
- Hopelessness or despair – “What’s the point?”
- Memory blanks – poor recall, difficulty staying present.

Behavioral Markers

- Withdrawal – avoids eye contact, social disengagement.
- Freeze – immobility, unable to take action even when needed.
- Collapse behaviors – lying down, curling up, “checked out.”
- Sleep – excessive or “shutdown” sleep that doesn’t restore.

Clinical Note for Therapists

Dorsal vagal activation is often misread as calm — but the body is not relaxed, it’s collapsed. Clients may present as “quiet” or “compliant,” but their nervous system is in shutdown, not safety. Look for the tell-tale signs: blank eyes, flat voice, limp gestures, and “I don’t care” language.

References

1. Thayer JF, Åhs F, Fredrikson M, Sollers JJ, Wager TD. A meta-analysis of heart rate variability and neuroimaging studies: Implications for heart rate variability as a marker of stress and health. *Neurosci Biobehav Rev.* 2012;36(2):747-756.
2. Levine P. In *An Unspoken Voice: How the Body Releases Trauma and Restores Goodness*. Berkeley: North Atlantic Books; 2015.
3. Mayer EA. The neurobiology of stress and gastrointestinal disease. *Gut.* 2011;60(2):307-317.
4. Porges SW. *The Pocket Guide to the Polyvagal Theory*. New York: Norton; 2017.


Understanding the ANS is the beginning. To bring this knowledge into the therapy room, we need ways to observe and track the nervous system in real time.

How do we know if a client is shifting into dorsal shutdown, stuck in sympathetic overdrive, or finding ventral vagal safety? The next page introduces the four key 'mirrors' of the ANS — tools that help us see and measure what's happening beneath the surface.

Part 3

How do we track the ANS?

To truly support regulation, we need ways to see the nervous system in action. These four aspects offer windows into how the autonomic nervous system is functioning. They serve as markers of balance and dysregulation, helping us read what the body is saying—often before words can.



Window of Tolerance

Mapping the Nervous System in Real Time

"Where the nervous system can do life without overwhelm or shutdown."

Most of us move in and out of our Window of Tolerance (WOT) every day. But when that window becomes narrow—through trauma, chronic stress, or years of overriding our body's signals—life feels harder to hold.

What a small window looks like in practice:

- You tire quickly and feel drained — there's simply less to give emotionally, creatively, and cognitively.
- You've spent years "pushing through" instead of listening to your body.
- Needs and boundaries have been ignored or overridden—often bottled up under layers of responsibility.
- Chronic stress, trauma, and even not living your truth can squeeze the window even tighter.

A narrow WOT doesn't mean the client is "broken." It means their system is working overtime to keep them safe. Our goal is not to stop activation — it's to gently widen the window so the system can flex and return to balance.

Clinical Insight

This is one of your most important tools.

Your entire therapy process will orbit around helping clients expand their window.

Before choosing any intervention, first ask:


"Is my client inside, above, or below their window of tolerance?"

Everyday WOT "situations":

The MOM...

The open-mouth-chewing-dad and the teenager...

The too much coffee...



Levine, Ogden and Siegal


What Is the Window of Tolerance?

Coined by Dr. Dan Siegel, the Window of Tolerance (WOT) refers to the optimal arousal zone where we can think clearly, connect relationally, feel emotions safely, and respond adaptively to challenges.

When we are **within this window**, the prefrontal cortex is online, and the autonomic nervous system is balanced.

Outside the Window...

When we leave the window due to real or perceived threat, we enter autonomic survival states:



The Healthy
Window Is
Flexible, Not Fixed

- A narrow window = less resilience to stress, more reactivity or collapse
- A widened window = greater emotional regulation, adaptability, and presence
- A dysregulated window = cycling between hyper- and hypo-states, often unconsciously


References

1. Porges SW. The polyvagal theory: Neurophysiological foundations of emotions, attachment, communication, and self-regulation. Norton; 2011.
2. Dana D. The Polyvagal Theory in Therapy. Norton; 2018.
3. Kline R. Clinical applications of the vagus nerve in trauma therapy. J Mind-Body Integration. 2020;4(1):15-22.


What Narrows the Window?

- Early trauma or attachment wounds
- Chronic stress or emotional suppression
- Sensory processing issues
- Neuroimmune dysregulation
- Unprocessed shock, grief, or fear


Increasing Your Window of Tolerance



Narrow Optimal Zone:
Limited Capacity to handle Stress



Wide Optimal Zone:
Resilient & Resourceful under Stress



Quick Reference - The Window Map

State	Description	Nervous System	Clinical Markers	Clinical Priority
Above WOT	Hyperarousal	Sympathetic	Anxiety, racing thoughts, muscle tension, hypervigilance	Contain arousal, rhythm, breath pacing
Inside WOT	Regulation	Ventral vagal	Calm focus, relational engagement, flexible emotion	Build reflection, insight, higher therapy work
Below WOT	Hypoarousal	Dorsal vagal	Numb, flat, dissociated, fatigue	Activate gently, sensory engagement, orienting

Clinical Insight - "State First, Content Second"

Therapist Reminder:

- Always observe state before exploring story.
- Clients outside their window:
 - Cannot fully process emotions
 - Cannot integrate insight
 - Might not engage in therapy fully
 - Might not progress
 - May dissociate or retraumatize if pushed cognitively

Self-Check &

- Do you track your client's window actively during sessions?
- How do you notice when a client is starting to shift out of their window?
- What are your own body cues when a client becomes dysregulated?

Clinical Applications:

- Therapists must assess where the client is in their window before choosing interventions.

Ask:

- Can they feel emotion without being overwhelmed?
- Are they able to stay in their body while speaking?
- Do they show signs of shutdown or flooding?

Clients outside their window cannot benefit fully from cognitive therapy or trauma processing

How to Use This in Practice:

Client is frozen → low affect → Hypoarousal → Begin with somatic awakening, gentle sensory input
Client is anxious → scattered → Hyperarousal → Use orienting, breath pacing, grounding tools
Client is present → aware → Window → Proceed with relational work, cognitive reflection

What We're Really Tracking:

The moment-to-moment shifts in autonomic state beneath the words.

The body reveals more than language

References

1. Siegel DJ. The Developing Mind: How Relationships and the Brain Interact to Shape Who We Are. Guilford Press; 2012.
2. Ogden P, Minton K, Pain C. Trauma and the Body: A Sensorimotor Approach to Psychotherapy. Norton; 2006.
3. van der Kolk BA. The Body Keeps the Score. Penguin; 2014.

Interoception

"The body's first language."

Interoception is the sense that tells us what's happening inside — the thud of the heart, the flutter in the stomach, the tightness in the chest before we can name it as fear, excitement, or dread. It's the body's constant stream of updates, whispering what it needs: more air, more rest, more safety.

When interoception is clear, we notice those signals early. We eat when we're hungry, we rest when we're tired, we pause when something feels "off." When it's dulled or disconnected, we push through warning signs, confuse fatigue with laziness, or call anxiety "just who I am."

If self-awareness was the house, interoception would be the foundation — quiet, unseen, but holding everything steady. When the foundation cracks, the whole structure feels unstable. When it's restored, we can build true awareness, regulation, and resilience on top of it.

Interoception: The Key to Noticing Dysregulation

What Is Interoception?

Interoception is the nervous system's ability to perceive internal body states—such as heart beat, breath rhythm, temperature, digestion, pain, and emotional arousal. It is your body's internal sensory system—the source of felt sense, self-awareness, and internal safety.

Why It Matters for Regulation

Clients cannot regulate what they cannot sense...

Interoception is what allows a person to:

- Notice early signs of anxiety (e.g. increased heart rate)
- Feel hunger or fullness
- Detect tension, heat, or overwhelm
- Recognize emotion as it emerges (before it escalates)

It forms the foundation of self-awareness and is central to trauma recovery, emotional regulation, and body-based therapy.

Interoception Links Brain to Body

Key brain regions involved in interoceptive processing include:

- Insula: Primary interoceptive hub
- Anterior cingulate cortex (ACC): Emotional regulation and error detection
- Thalamus: Sensory relay system
- Brainstem and vagus nerve: Real-time body monitoring

These structures interpret signals from the body and determine whether we feel safe or threatened.

When Interoception Is Disrupted...

- The client may not notice their breath speeding up
- They may misinterpret internal cues (e.g. hunger as anxiety)
- Emotions may build unnoticed until they explode or collapse
- They may feel "numb," disembodied, or alienated from their body

Interoceptive deficits are common in:

- Trauma survivors
- Anxiety disorders
- Depression
- Neurodivergent individuals (e.g. ASD, ADHD)
- Somatic dissociation

Clinical Insight:

"Interoception is the bridge between survival and self-regulation."

When clients learn to feel their body cues without fear, they begin to reclaim the ability to regulate in real time.

Awareness and regulation help improve interoception.

References

1. Quadt L, Critchley HD, Garfinkel SN. The neurobiology of interoception in health and disease. *Ann NY Acad Sci.* 2018;1428(1):112–28.
2. Nord CL, Garfinkel SN. Interoceptive pathways to mental health: linking body and brain. *Trends Neurosci.* 2022;45(6):447–59.

Clinical Insight

Interoception is what allows your clients to notice their dysregulation before it spirals. Without interoception, clients stay disconnected from their bodies.

Your goal is not just to calm — it's to help clients feel safely again.

Quick Reference Clinical Markers of Interoceptive Disruption

Symptom	What It Reveals
"I don't feel anything"	Dorsal collapse; interoceptive blunting
"I notice my heart races but too late"	Delayed interoceptive signal recognition
"I confuse hunger with anxiety"	Signal misinterpretation
Sudden emotional flooding	Poor early detection of rising arousal
Panic attacks "out of nowhere"	Missed early bodily cues
Body feels "numb" or "alien"	Somatic dissociation, interoceptive shutdown

Clinical Practice:

Use simple prompts to begin rebuilding interoceptive awareness:

- "Notice your breath — is it moving in your chest or belly?"
- "Where in your body feels tight right now?"
- "If this sensation had a color, what would it be?"
- "Does that sensation change as we sit together?"

- Do you regularly help your clients track internal body cues?
- Do you normalize noticing physical sensations as part of healing?
- What interoceptive signals do you feel when clients become activated?

Neuroception & Safety Cues

"The body's silent question: Am I safe?"

Neuroception is the nervous system's surveillance — a wordless, constant scanning of the world, long before thought or logic. It notices the micro-tensions in a stranger's face, the tone in a voice, the shift of a shadow across a room. All of it is measured not with reason, but with instinct. When neuroception senses safety, the body softens. Breath steadies. The world feels open. When it detects danger — even if we can't name why — muscles tighten, the jaw clenches, and the heart prepares for what might come.

If interoception is the foundation of self-awareness, neuroception is the lock on the door — quietly deciding what enters, what stays outside, and how much of ourselves we can bring into the world at any moment.

Neuroception in Action

- When you stand in front of a toxic coworker, your body knows before you do — shoulders tense, your breath goes shallow, and your heart rate rises, even as you smile politely.
- When you see something traumatic on the news, your stomach knots and your throat tightens. You may not realize it, but your nervous system just filed that scene under "threat."
- When you hear a sudden loud noise in a parking lot, your muscles coil before your mind even processes what happened.
- When a stranger's tone is warm and soft, your body unclenches; your eyes lift, you lean in — the world feels safer without you even thinking about it.
- When you walk into an unfamiliar room, you instinctively scan faces, doorways, and exits. You might call it "reading the room," but your body has already read it for you.

Smile...

Neuroception & Safety Cues: How the Body Decides Who to Trust

What Is Neuroception?

Coined by Dr. Stephen Porges, neuroception refers to the nervous system's subconscious ability to detect safety, danger, or life threat — without involving conscious thought.

Before we feel, think, or act, the body asks:

- “Am I safe, or not?”

This decision determines whether the body opens toward connection or moves into survival responses like fight, flight, or freeze.

“It’s not what we think. It’s what our body senses.”

Neuroception vs. Perception

Perception

Cognitive, conscious

Happens in the thinking brain

Based on logic, memory, language

“I know I’m safe.”

Neuroception

Subcortical, unconscious

Happens in the body and brainstem

Based on facial tone, voice prosody, posture, context

“I feel safe.” (or not)

What the Body Scans For:

- Facial expressions (through mirror neurons)
- Tone of voice (prosody)
- Posture and movement of others
- Sensory environment (light, sound, texture, smell)
- Internal cues (heart rate, gut tension, breath pattern)

These are interpreted instantly to decide whether to activate ventral vagal connection, sympathetic mobilization, or dorsal vagal shutdown.

When Neuroception Goes “Off”

In trauma survivors, neuroception may misread safe cues as dangerous:

- A kind touch feels threatening
- Eye contact is intolerable
- Silence feels like abandonment
- Rest triggers anxiety

This faulty neuroception is not defiance — it’s the body’s way of saying:

“This has hurt me before. I can’t trust it yet.”

Clinical Insight:

To restore healthy regulation, the therapist must become a source of co-regulated safety. Your tone, facial expression, movement, and pace matter more than your words.

Clinical Anchor:

Neuroception is not a cognitive process. It is the body's surveillance system, scanning for cues of danger or safety every moment.

If neuroception says "unsafe," no amount of logic will help.

Your presence and environment must speak "safe" at the level of the body.

Quick Reference Insert - Neuroception Pathways

Neuroception happens through multiple sensory channels. Here's how the body takes in cues — and how you'll see it play out in your clients

Input Type	System Pathway	Clinical Signs
Visual (facial expression, eye contact)	Superior colliculus → amygdala → vagus	Client looks away, freezes, scans for threat
Auditory (tone, prosody)	Middle ear → brainstem → vagus	Client responds to tone more than content
Postural (body movement, proximity)	Proprioception → cerebellum → insula	Client tenses or moves away reflexively
Internal (gut, heart, interoception)	Afferent vagus → brainstem → insula	Client reports "bad feeling" or gut sense

Therapist Tools for Regulating Neuroception:

- Soften your face — the client reads your tension first.
- Speak with warm, slow, prosodic tone.
- Sit at a slight angle (not head-on).
- Reduce visual clutter and harsh lighting.
- Match their breath rhythm nonverbally before guiding them.
- Normalize body-based safety talk:

"Your body is wise — let's follow it."

References

1. Porges SW. The polyvagal perspective. *Biol Psychol.* 2007;74(2):116–43.
2. Porges SW. The polyvagal theory: neurophysiological foundations of emotions, attachment, communication, self-regulation. New York: Norton; 2011.
3. Dana D. The polyvagal theory in therapy: engaging the rhythm of regulation. New York: Norton; 2018.
4. Kleckner IR, Zhang J, Touroutoglou A, Chanes L, Xia C, Simmons WK, et al. Evidence for a large-scale brain system supporting allostasis and interoception in humans. *Nat Hum Behav.* 2017;1(5):69.
5. Critchley HD, Harrison NA. Visceral influences on brain and behavior. *Neuron.* 2013;77(4):624–38.
6. Stephan KE, Manjaly ZM, Mathys CD, Weber LAE, Paliwal S, Gard T, et al. Allostatic self-efficacy: a metacognitive theory of dyshomeostasis-induced fatigue and depression. *Front Hum Neurosci.* 2016;10:550.

Allostatic Load

The body remembers. Even when the mind tries to move on.

The Weight We Can't See

Your nervous system was never built to hold the storm forever.

It was built to let the rain pass, the sun return, the air clear.

Allostatic load is what happens when the storm doesn't leave —
when the stress of yesterday, last month, last year
never fully drains from the body.

Think of your body like a roof in heavy rain.

A little rain is fine — it beads up, slides off,
the roof holds steady.

But when the rain becomes a downpour,
and the gutters are clogged with old leaves,
water pools, weight builds, cracks form.

That's allostatic load:

the invisible weight of too many "just one more days" of stress.

It's the reason someone can "push through" for years —
smiling, working, surviving —
and then one day their body says, "no more."

Like a bridge carrying too much traffic,
or a credit card that's been swiped too many times,
the system defaults. The price must be paid.

Allostatic load is not weakness.

It's the biology of accumulation —
the way cortisol drips like a slow leak,
how inflammation simmers quietly,
how sleep frays and repair is delayed.

It's the shoulders that never drop,
the stomach that knots before every meeting,
the sigh you don't even know you're holding.

If self-regulation is a savings account, allostatic load is the overdraft.

Every deposit of rest, safety, and joy matters —
but so does every withdrawal.

Your body is keeping score,
but it's also waiting for balance.

For the gutters to be cleared,
for the rain to finally run free.

Allostatic Load & Trauma Looping: When the System Gets Stuck in Survival

What Is Allostatic Load?

Allostasis is the body's ability to achieve stability through change. It's how we adapt to life's demands — adjusting heart rate, hormones, attention, and energy to meet what's needed. But when stress is chronic or overwhelming, the system pays a cost.

This cumulative burden is called allostatic load — the “wear and tear” on the nervous system and body over time.

Allostatic Load vs Homeostasis

Homeostasis = Maintaining a stable internal state (e.g. body temp)

Allostasis = Actively shifting internal resources in response to stress

Allostatic Load = The cost of repeated adaptation — often invisible, until breakdown occurs

What Increases Allostatic Load?

- Early childhood trauma or neglect
- Persistent relational conflict
- Chronic illness or sleep deprivation
- Financial insecurity, overwork, burnout
- Unprocessed grief, fear, shame, or hypervigilance
- Even thinking about past danger (rumination)

The body doesn't distinguish between past and present danger — if the memory still has charge, the body pays the price.

Trauma Looping: When the System Gets “Stuck”

When the nervous system is repeatedly activated without completion or repair, it becomes pattern-locked in protective states.

“What was once adaptive becomes automatic.”

This can result in:

- Persistent sympathetic arousal (anxiety, GI issues, hypervigilance)
- Dorsal collapse (depression, shutdown, chronic fatigue)
- Oscillating between the two with no recovery
- Diminished ability to return to the window of tolerance

Clinical Insight: Allostatic Load and the Shrinking Window of Tolerance

Definition Reminder:

Allostatic load is the cumulative physiological cost of chronic exposure to stress.

It's not just about having stress — it's about what happens when the body must stay in survival mode for too long without recovery.

How Allostatic Load Shrinks the Window of Tolerance:

Allostatic Process	Effect on Nervous System	Result on WOT
Chronic cortisol & adrenaline activation	Sensitizes amygdala, inhibits PFC	Faster triggers into hyperarousal
Vagal withdrawal & shutdown patterns	Suppresses interoception, co-regulation	Slower recovery → prolonged hypoarousal
Impaired feedback loops (HPA axis, immune, sleep)	Breaks natural recovery mechanisms	Baseline moves closer to dysregulation
Emotional suppression / over-functioning	Keeps system in tension	Narrower WOT with false "calm" masking tension

Clinical Markers of Allostatic Overload:

- Hyperreactivity to minor stressors ("exploding over small things")
- Quick collapse into shutdown, fatigue, or disconnection
- Physical complaints (chronic pain, gut issues, hormonal shifts)
- Difficulty regulating even with familiar strategies
- Flattened affect or "numb busyness"
- Inability to rest even when safe
- Poor sleep, brain fog, immune flares

Bottom Line:

Allostatic load is invisible weight. It compresses the Window of Tolerance not because of current trauma — but because the body never got a break from past survival. Your work is to offer:

- Restoring rhythm
- Co-regulated pause
- Space to feel without overflow

And in time, the window widens again.

Clinical Markers of High Allostatic Load

Domain	Symptom
Physical	Chronic pain, inflammation, sleep issues, fatigue
Cognitive	Brain fog, memory loss, inability to focus
Emotional	Rewinding past events, shutdown, irritability
Behavioral	Withdrawal, compulsive coping (e.g., overworking, bingeing)

NOTE: the higher the allostatic load - the smaller the window of tolerance



What This Means for Therapy

High allostatic load clients cannot sustain insight-driven therapy

Their system must first experience regulation from the inside out

Repetition of survival experiences wires the body for defense

Repetition of safety experiences rewires it for resilience

References

1. McEwen BS. Protective and damaging effects of stress mediators. *N Engl J Med.* 1998;338(3):171-179.
(Seminal paper introducing the concept of allostatic load.)
2. McEwen BS, Seeman T. Protective and damaging effects of mediators of stress. *Ann N Y Acad Sci.* 1999;896:30-47.
3. Van der Kolk BA. *The Body Keeps the Score.* New York: Viking; 2014.
(Links chronic stress and trauma to physiological imprinting.)
4. Porges SW. *The Pocket Guide to Polyvagal Theory.* New York: Norton; 2017.
5. Juster RP, McEwen BS, Lupien SJ. Allostatic load biomarkers of chronic stress and impact on health and cognition. *Neurosci Biobehav Rev.* 2010;35(1):2-16.
(Comprehensive overview of biomarkers like cortisol, CRP, DHEA, metabolic markers.)
6. Sterling P. Allostasis: a model of predictive regulation. *Physiol Behav.* 2012;106(1):5-15.
(Deepens understanding of allostasis as a predictive, energy-balancing system.)
7. Danese A, McEwen BS. Adverse childhood experiences, allostasis, allostatic load, and age-related disease. *Physiol Behav.* 2012;106(1):29-39.
(Links early trauma to long-term allostatic burden and disease risk.)
8. Karatsoreos IN, McEwen BS. Psychobiological allostasis: resistance, resilience, and vulnerability. *Trends Cogn Sci.* 2011;15(12):576-584.
(Important for therapists: introduces resilience vs. vulnerability in stress systems.)


Part 4

Scientific Biomarkers

While the body tells its story through sensations, behaviors, and patterns, some signals can also be measured. Biomarkers are the body's fingerprints of regulation — tangible data points that reflect what's happening deep in the autonomic nervous system.

They don't replace observation and intuition — they complement them. Just as a stethoscope doesn't replace listening to a patient's words, biomarkers don't replace listening to the body. But they give us an additional lens — an objective one — to see what the nervous system might not say aloud.

These two biomarkers — HRV and heart-brain coherence — have become powerful tools for clinicians and researchers alike. They translate the language of the autonomic nervous system into data we can see, track, and respond to.



Coherence

The Healing Rhythm Beneath Words

The body is built on rhythm.

Your heart beats in waves, your lungs rise and fall like tides, your brain hums with electric pulses.

When life is calm, these rhythms flow in harmony — heart, breath, hormones, and thoughts weaving together like instruments in the same song.

But when stress takes over, the rhythms shatter.

The heart pounds erratically, breath becomes shallow, brainwaves turn jagged — every system scrambling to survive.

Coherence is the return to harmony.

It's when the heart settles into a smooth, wave-like rhythm, and the breath falls into step with it.

Hormones rebalance — cortisol and adrenaline ebb, oxytocin and DHEA rise.

Inflammation quiets. The immune system wakes.

Brainwaves slow into alpha and theta, opening the door for clarity, focus, and emotional integration.

Coherence is not just calm — it's the body's systems learning to speak the same language again.

It is the rhythm that allows healing to begin, not through force, but through flow.

If the body were an orchestra, coherence is the conductor —

lifting the baton to bring heart, breath, and mind back into one song.

Coherence: The Healing Rhythm Beneath Words

“The nervous system doesn’t heal through language. It heals through rhythm.”

What Is Coherence?

According to the HeartMath Institute:

“Coherence is a state in which the heart, mind, and emotions are in energetic alignment and cooperation.”

Physiologically, coherence refers to a synchronized, sine-wave pattern in heart rate variability (HRV). It represents optimal function across multiple body systems, including:

- Cardiovascular
- Respiratory
- Nervous system
- Endocrine
- Immune
- Cognitive and emotional integration

In coherence, there is efficiency, flow, and balance — a state where the body’s systems are working with one another, not in conflict.

Coherence and the Autonomic Nervous System (ANS)

The ANS regulates heart rate, breath, digestion, and threat response.

In dysregulation:

- Systems become disconnected or overactive
- Transitions between sympathetic and parasympathetic states become rigid
- Clients remain “stuck” in survival responses

In coherence:

- Vagal tone increases, improving parasympathetic flexibility
- Transitions between states become smoother
- The body returns more easily to baseline
- The window of tolerance expands — meaning the client can tolerate more intensity without flipping into shutdown or overwhelm

Coherence is the felt sense of being in your window of tolerance.

What Happens in the Body During Coherence?


System	Change in Coherence
Cardiovascular	Heart rate rhythms become smooth and wave-like (HRV increases)
Respiratory	Breathing deepens, slows, and syncs with heartbeat (RSA increases)
Endocrine	Cortisol and adrenaline decrease, oxytocin and DHEA increase
Immune	Inflammation markers drop; immune resilience improves
Cognitive	Frontal cortex activity increases (clarity, focus, memory)
Emotional	Emotion regulation improves; amygdala reactivity decreases
Neurological	Brainwaves shift from high-beta (stress) to alpha/theta (flow, safety)

Clinical insight

Coherence breathing keeps the nervous system within the window of tolerance, allowing access to the prefrontal cortex — so the client can stay regulated and reflective.

HRV Patterns (Coherent vs. Incoherent)

Heart Rhythms (Heart Rate Variability)



Clinical insight

Coherence is not relaxation it is a measurable state where the heart rhythm, breathing rate, and brain function enter a synchronized, harmonious rhythm. It reflects **a state of regulation, presence, and nervous system adaptability** – not just “calm.”

✓ High coherence = flexible, resilient

✗ Low coherence = reactive, fragmented

AND

Clinical Note:

HRV (Heart Rate Variability) is the measurement — coherence is the pattern.

You can have high HRV but low coherence.

We want both: flexible variability and a stable rhythm.

Brain States and Brainwaves in Coherence

- High beta (25+ Hz) – Linked to stress, anxiety, overdrive
- Alpha (8–12 Hz) – Relaxed focus, present attention
- Theta (4–7 Hz) – Access to deep emotion, imagination, memory
- Delta (<4 Hz) – Deep sleep, restoration

During coherence:

- The brain shifts out of high beta (chaos)
- Alpha and theta rise — creating space for emotional integration, healing, and creativity

Emotion and Coherence

HeartMath has shown that:

- Positive emotions (gratitude, compassion, appreciation) increase coherence
- Negative emotions (frustration, fear, shame) disrupt coherence, even when breathing remains slow

This proves:

- Coherence is not just physical — it's emotional alignment
- The quality of emotion matters as much as breath or posture
- This is why therapeutic attunement and emotional safety are so critical
- “Coherence isn't just calm — it's the rhythm of safety, gratitude, and presence.”

Clinical Application

- Use coherence as the baseline goal for regulation, not relaxation
- Observe facial tone, breath rhythm, and heart-emotion congruence
- Teach clients that emotional resonance is part of their healing rhythm
- Encourage use of coherence tracking tools (see next page)

Therapist Coherence = Client Regulation

- Mirror neurons + neuroception mean your coherence entrains the client
- In sessions, focus on becoming a coherent nervous system
- Clients will match your breath, tone, pace, and facial affect without needing explanation

“The therapist's coherence is the intervention.”

References

1. McCraty R, Zayas M. Cardiac coherence, self-regulation, autonomic stability, and psychosocial well-being. *Front Psychol.* 2014;5:1090.
2. Lehrer P, Vaschillo E, Vaschillo B. Resonant frequency biofeedback training to increase cardiac variability: rationale and manual for training. *Appl Psychophysiol Biofeedback.* 2000;25(3):177–191.
3. Porges SW. *The Polyvagal Theory: Neurophysiological Foundations of Emotions, Attachment, Communication, and Self-Regulation.* New York: W.W. Norton; 2011.
4. Dana D. *The Polyvagal Theory in Therapy: Engaging the Rhythm of Regulation.* New York: W.W. Norton; 2018.
5. Buteyko KP. *The Buteyko Method: Breathing for Health.* Moscow: Buteyko Clinic; 1991.
6. HeartMath Institute. *Science of the Heart: Exploring the Role of the Heart in Human Performance, Vol. 2.* Boulder Creek, CA: HeartMath Institute; 2016.

Clinical Insight: Brainwaves Reflect Nervous System State

Our brain doesn't operate in isolation — its electrical activity mirrors what's happening in the autonomic nervous system.

- High Beta (fast, spiky) → Often linked to sympathetic arousal — anxiety, overthinking, stress
- Low Alpha or Theta flooding → Associated with dorsal vagal shutdown, daydreaming, dissociation, or freeze
- Balanced Alpha → Reflects ventral vagal tone, calm focus, ideal learning or therapeutic state

Heart Rate Variability

What Your Heartbeat Reveals About Your Life

Heart Rate Variability: What Your Heartbeat Reveals About Your Life

“Your heartbeat is not a clock. It’s a conversation between your body and your brain.”

What Is Heart Rate Variability (HRV)?

- Heart Rate Variability (HRV) is the variation in time between heartbeats — not the heart rate itself, but the rhythm between beats. A healthy, regulated heart does not beat evenly like a metronome.

Instead, it responds to:

- Emotions
- Breath
- Stress
- Hormonal changes
- Safety signals

*This variability reflects how flexible and responsive your nervous system is.
HRV is the fingerprint of your autonomic nervous system.*

Why HRV Matters

HRV is considered one of the most reliable biomarkers of nervous system regulation and overall health. It reflects:

- Vagal tone (strength of the parasympathetic system)
- Stress resilience
- Emotional regulation capacity
- Adaptability of the body to internal and external demands

Low HRV is not just about poor fitness — it’s often about nervous system overload, trauma, inflammation, or shutdown.

What Influences HRV?

Breathing -----Slow, nasal breathing increases HRV; mouth breathing or over-breathing reduces it

Emotions -----Gratitude and compassion raise HRV; frustration and fear lower it

Posture & Movement --Slumped posture and immobility reduce HRV; gentle rhythmical movement supports it

Sleep & Recovery -----Poor sleep reduces HRV; deep rest restores it

Trauma -----Past trauma reduces baseline HRV, especially in dorsal vagal states

Clinical insight -

HRV is the most reliable non-invasive measure of vagal tone — the health of the vagus nerve and its ability to regulate your system.

➤ A higher vagal tone = quicker recovery from stress and better emotional regulation.

What Does HRV Reveal?

HRV Pattern	What It May Indicate
High & smooth	Resilient regulation, strong vagal tone
Low & flat	Chronic stress, burnout, vagal suppression
Erratic	Anxiety, trauma response, poor interoceptive stability
High but incoherent	Over-breathing or sympathetic hyperarousal with some control

HRV and the Nervous System

- The vagus nerve controls heart rhythm via the sinoatrial (SA) node
- Strong vagal tone = heart can quickly shift states and return to baseline
- Weak vagal tone = stuck patterns, poor recovery, survival state looping

“If the heart can’t dance, the brain can’t think clearly.”

Clinical Insight: How to Use HRV Clinically

- Track HRV over time, not just in the moment
- Watch for daily trends: is it rising with regulation practice?
- Combine HRV with subjective observation: breath pattern, posture, speech, engagement
- Teach clients the link between how they breathe, how they feel, and how their heart responds

Note: HRV biofeedback provides a mirror to the nervous system — clients learn to recognize and shift state in real time.


References

1. McCraty R, Shaffer F. Heart rate variability: new perspectives on physiological mechanisms, assessment of self-regulatory capacity, and health risk. *Glob Adv Health Med*. 2015;4(1):46–61.
2. Lehrer P, Vaschillo E, Vaschillo B. Resonant frequency training: clinical applications and physiological mechanisms. *Appl Psychophysiol Biofeedback*. 2000;25(3):177–191.
3. Kim H-G, Cheon E-J, Bai D-S, Lee YH, Koo BH. Autonomic dysfunction in trauma: heart rate variability as a biomarker of trauma-related dysregulation. *Psychiatry Investig*. 2018;15(3):232–239.
4. HeartMath Institute. *Science of the Heart: Exploring the Role of the Heart in Human Performance*, Vol. 2. Boulder Creek, CA: HeartMath Institute; 2016.
5. Thayer JF, Lane RD. A model of neurovisceral integration in emotion regulation and dysregulation. *J Affect Disord*. 2000;61(3):201–216.
6. Shaffer F, Ginsberg JP. An overview of heart rate variability metrics and norms. *Front Public Health*. 2017;5:258.

Part 5

How the Nervous System Learns Regulation: Loops, Mirrors & Active Engagement

The nervous system is not a static wiring — it is sculpted daily by patterns (neuro-loops), reflections from others (mirror neurons), and the actions we take (active participation).



Neuro-Looping

How the Brain Repeats What It Survives

"We don't just remember trauma. We repeat its rhythm."

The nervous system is a storyteller...

It doesn't just hold memories — it replays them.

When a trigger hits, your body can pull out an old script and perform it like it's opening night.

The brain fires. The body reacts. The behavior confirms.

And the loop tightens — even when the danger is long gone.

This is not weakness. It's wiring.

The nervous system learned what kept you alive. And like any loyal student, it keeps repeating the lesson — even when the exam is long over.

But here's the hope: loops can be rewritten.

New experiences of safety, rhythm, and co-regulation create new scripts.

Therapy doesn't just interrupt old loops — it teaches the nervous system a different ending.

Neuro-looping refers to the cyclical pattern...

What Is Neuro-Looping?

Neuro-looping refers to the cyclical pattern in which the brain, body, and nervous system replay survival responses — even when the original danger is long gone. It's not just habit.

It's a biological loop built through:

- Repetition of stress signals
- Reinforcement through body memory
- Neural wiring that prefers what is familiar, even when it's dysfunctional

“What fires together, wires together — even if what's firing is fear.”

How a Loop Forms

1. **Trigger:** A stimulus (external or internal) activates a memory or body sensation
2. **Perception:** Neuroception misinterprets the present through the lens of past pain
3. **Response:** The body enters a survival state (fight, flight, freeze, fawn)
4. **Reinforcement:** The nervous system adapts around this state — rewiring to expect it
5. **Repetition:** With each loop, the neural pathway strengthens, becoming the default

This is how the body wires in survival: trigger → perception → response → reinforcement → repetition.

This is why regulation isn't optional — it's how we open a new loop before the old one runs again

Common Neuro-Loops in Clients

Loop Type	Description	Nervous System Profile
Anxiety-Overdrive Loop	Hyperarousal → overwhelm → avoidance → shame → more anxiety	Sympathetic dominance
Freeze-Collapse Loop	Inactivity → shame → hopelessness → more shutdown	Dorsal vagal dominance
Fawn-Exhaustion Loop	People-pleasing → burnout → panic → perform again	Mixed state (fawn = parasympathetic + sympathetic)
Pain-Fear Loop	Pain → fear of movement → immobility → more pain	Allostatic overload
Cognition-Overcontrol Loop	Overthinking → fear of emotion → disembodiment → more overthinking	Prefrontal overactivation with vagal disconnection

Why It Matters

Neuro-looping explains why clients:

- Keep ending up “back where they started”
- Feel confused about why they “sabotage” progress
- Say “I understand it, but I can’t change it”

These are not psychological failures — they’re physiological loops.

Clinical Strategy:

Map the Loop Before You Interrupt It

Therapists must become loop mappers, not just problem solvers.

Ask:

What is the first spark in this loop?

Where does the client leave their window of tolerance?

What behavior is their nervous system using to protect, not sabotage?

Clinical Insight

A neuro-loop is a repeated firing pattern between brain regions, body sensations, thoughts, and behaviors — originally wired in response to survival, then replayed automatically, even when the original danger is gone.

“The nervous system doesn’t repeat what’s logical — it repeats what helped us survive.”

This is how maladaptive loops form: the body reacts → brain justifies → behavior confirms → brain wires it in.

Practical Examples of Common Neuro-Loops:

Loop 1: I am a failure

Trigger: Criticism

Sensation: Gut tension, heat in face

Thought: “I’m a failure”

Behavior: Withdraw, self-blame, perfectionism

→ Reinforced by: relief from avoiding more shame

Loop 2: I can’t cope

Trigger: Performance pressure

Sensation: Racing heart, blank mind

Thought: “I can’t cope”

Behavior: Collapse, forget, dissociate

→ Reinforced by: no consequences if you shut down

Loop 3: People Please

Trigger: Conflict or disapproval

Sensation: Tight throat, shallow breath

Thought: “If I upset them, I’ll be rejected”

Behavior: Agree, smile, suppress

→ Reinforced by: short-term safety, no rejection

If you suspect a neuro-loop is active, begin by gently slowing the session and inviting the client to track their body’s automatic response before analyzing thoughts or behavior. Say something like, “Let’s pause — what do you notice in your body right now?” This interrupts the default survival pathway. Once the client is present, help them identify the loop’s trigger, sensation, thought, and action pattern, then offer a bottom-up regulation cue (such as a breath reset, orienting movement, or grounding touch) to disrupt the loop at its origin point — the body. This opens space for a new, regulated pattern to form before the old loop replays.

References

1. LeDoux JE. The Emotional Brain: The Mysterious Underpinnings of Emotional Life. Simon & Schuster; 1996.
2. van der Kolk BA. The Body Keeps the Score: Brain, Mind, and Body in the Healing of Trauma. Viking; 2014.
3. Porges SW. The Pocket Guide to Polyvagal Theory: The Transformative Power of Feeling Safe. W.W. Norton; 2017.
4. Hebb DO. The Organization of Behavior: A Neuropsychological Theory. Wiley; 1949.
5. Damasio AR. The Feeling of What Happens: Body and Emotion in the Making of Consciousness. Harcourt; 1999.
6. Schore AN. Affect Regulation and the Origin of the Self: The Neurobiology of Emotional Development. Routledge; 2016.
7. Siegel DJ. The Developing Mind: How Relationships and the Brain Interact to Shape Who We Are. Guilford Press; 2020.
8. Kolb B, Gibb R. Brain plasticity and behaviour in the developing brain. J Can Acad Child Adolesc Psychiatry. 2011;20(4):265–276.
9. Sapolsky RM. Why Zebras Don’t Get Ulcers. Holt; 2004.
10. McEwen BS. Protective and damaging effects of stress mediators. N Engl J Med. 1998;338(3):171–179.

Mirror Neurons

Mirror Neurons and the Relational Brain: Why Your Calm Regulates Others

"The body learns safety by watching yours."

What Are Mirror Neurons?

Mirror neurons are a special class of brain cells that activate both when:

- We perform an action, and
- We observe someone else performing that same action.

Discovered in the 1990s by researchers studying motor activity in monkeys, these neurons explain how we learn through imitation, experience empathy, and resonate emotionally with others — even without words.

The Nervous System Learns by Watching

Your client's brain is constantly scanning you — not just for what you say, but for:

- The tone of your voice
- The softness or tension in your face
- The depth of your breath
- The stillness or speed of your movements
- The presence in your eyes

These cues are processed by their mirror neuron system and either:

- Reinforce a sense of safety, or
- Confirm a need for defense

"When you breathe slower, they do too — often without realizing it."

When Mirror Neurons Have Been Trained for Danger

In trauma survivors or neglected children:

- Mirror neurons may be over-attuned to threat (e.g. mirroring anger, chaos)
- They may be underdeveloped (difficulty reading cues or empathizing)
- Eye contact or presence may feel unbearable, not bonding

"If the only face I learned from was unpredictable, every face feels dangerous now."

Therapist Clinical Application

- *Use your own nervous system as the intervention*
- *Model regulated breath, rhythm, and emotional presence*
- *Allow the client's mirror system to entrain to your safety — not your words*

This is neuroplasticity through embodiment.

Clinical tip:

- *Begin the session with 30 seconds of co-regulated breath*
- *Ask: "Can you feel your body noticing that I'm calm right now?"*
- *Reflect: "You didn't have to do anything. Your nervous system picked it up."*

References

- Rizzolatti G, Craighero L. The mirror-neuron system. *Annu Rev Neurosci.* 2004;27:169-192.
- Schore AN. *Affect Regulation and the Origin of the Self.* Lawrence Erlbaum; 2003.
- Porges SW. *Polyvagal Theory.* W.W. Norton; 2011.
- Dana D. *Polyvagal Theory in Therapy.* W.W. Norton; 2018.
- Cozolino L. *The Neuroscience of Psychotherapy.* W.W. Norton; 2017.

Clinical Insight:

- You are never neutral in the room.
- Your client's nervous system is reading yours — continuously.
- Mirror neurons and autonomic cues form a two-way loop.
- What you regulate in yourself, you regulate in them.

Quick Reference - Mirror Neurons in Therapy

Brain Area	Function	Clinical Relevance
Premotor Cortex	Action mirroring	Client mirrors therapist breath, posture, pace
Inferior Parietal Lobule	Intention recognition	Client senses therapist's emotional safety or tension
Insula	Empathic awareness	Shared affect; resonance or overwhelm
ACC	Emotional processing	Co-feeling → co-regulation loop builds or breaks

Active Participation in Therapy: Overview

Active Participation in Therapy: Overview

Definition:

Active participation refers to the client's voluntary, intentional involvement in:

Choosing goals

Engaging in activities

Reflecting on experience

Taking responsibility for aspects of their recovery or adaptation

It's a dynamic, co-created process — not passive treatment.

Why It Matters (Neuroscience Integration):

Neuroplasticity requires active engagement. The brain doesn't rewire through passive input alone — it needs intentional use, movement, choice, and reflection.

Nervous system regulation is embodied. Clients must feel safe enough to participate — and their participation deepens regulation.

Motor learning, sensory integration, and executive function all improve through active use, not observation alone.

When Clients Can't Actively Participate:

This is often a sign of dysregulation or neurophysiological shutdown. For example:

A child who "zones out" during fine motor tasks

An adult client who avoids social engagement, is fatigued, or says "I don't care"

A trauma client who cannot initiate movement or choices

In these cases, regulation becomes a precondition for participation.

Clinical Example:

A client with a stroke may not initiate movement in a hemiparetic arm. OT must assess:

Is this learned non-use, or dorsal vagal shutdown?

When regulation is restored (e.g. through co-regulated movement, breath, or rhythm), active participation re-emerges.

Nervous System Link

State	Participation Style	Therapist Response
Ventral vagal (regulated)	Engaged, curious, willing to try	Use co-creation, exploration
Sympathetic (fight/flight)	Anxious, over-achieving, or avoidant	Use rhythm, containment, orienting
Dorsal vagal (freeze)	Withdrawn, disengaged, passive	Use bottom-up sensory or movement co-regulation

Clinical Insight:

- Dysregulation blocks active participation.
- Participation isn't just compliance — it's internal ownership.
- Your task is to create enough safety for true participation to emerge.
- If the client is compliant but emotionally absent → they're not yet regulated.

Quick Reference - Participation Stages Cheat Sheet

Participation Stage	Client Behavior	Nervous System Profile	Clinical Focus
Survival / Withdrawal	Disconnected, flat, disengaged	Dorsal collapse	Safety, co-regulation, sensory awakening
Passive Compliance	Does tasks mechanically	Mixed dorsal / low ventral	Gentle rhythm, relational presence
Task-Oriented Activation	Goal-focused but emotionally guarded	Sympathetic-ventral blend	Contain perfectionism, maintain rhythm
Relational Engagement	Engaged, emotionally present	Ventral vagal activation	Expand reflection, relational safety
Integrated Participation	Fully present, flexible	Flexible regulation	Identity repair, meaningful occupation

Self-Check:

- In your own sessions:
 - Do you sometimes reward clients for compliance rather than genuine presence?
 - How do you track your client's participation level non-verbally?
 - How does your own nervous system feel sitting with a compliant but disengaged client?

Key Theories That Align:

- **MOHO** (Model of Human Occupation): Emphasizes volition, habituation, and performance — participation requires motivation and identity integration.
- **PEO Model** (Person-Environment-Occupation): Participation is where person and occupation meet in context. If the person is dysregulated, the “fit” collapses.
- **Sensory Integration Theory (Ayres)**: Participation is shaped by sensory processing and modulation — active engagement emerges from sensory safety.

Active Participation Requires Regulation: An OT Perspective

“Participation is not just doing. It is being present, engaged, and embodied in the task.”

Why Participation Matters

In occupational therapy, active participation is a foundational principle.

It refers to the client’s intentional involvement in therapeutic processes — through movement, choice-making, emotional presence, and meaningful engagement.

Participation is not passive compliance.

It involves:

Motivation (volition)

Habits and roles (habituation)

Sensory-motor readiness (performance capacity)

A regulated nervous system that allows for engagement

The Model of Human Occupation (MOHO) reminds us that true therapeutic change begins when the client is an active agent in their recovery — not just a recipient of intervention.

Why Regulation Comes First

Active participation requires access to the prefrontal cortex, the center of decision-making, planning, and reflection.

But in a dysregulated state (e.g. survival response, trauma activation), the body bypasses higher brain areas to prioritize safety.

When regulation is missing, participation isn’t resistance — it’s impossible.

Bottom Line:

Active participation begins with physiological safety.

Without regulation, there is no engagement — only survival.

Let this
sync

Level	Description	Nervous System State	Clinical Example
1. Survival Withdrawal	The client is disconnected from life. There is minimal or no engagement. Tasks feel unsafe. The world is too much.	Dorsal Vagal Shutdown	Flat affect, no eye contact, can't initiate action, avoids all responsibility
2. Passive Compliance	The client does what is asked, but without internal motivation. There's no felt meaning or drive. "Going through the motions."	Mixed state: low ventral, high dorsal	Does therapy homework mechanically, shows up but disengaged, avoids emotional topics
3. Task-Oriented Activation	The client begins to engage more purposefully with structured activities. Still avoids vulnerability or emotion, but begins movement toward life.	Sympathetic-Ventral Blend	Starts setting goals, completes tasks, becomes performance-focused but still emotionally guarded
4. Relational Engagement	The client initiates and maintains engagement. Emotional nuance returns. Begins showing curiosity, play, purpose.	Ventral Vagal Activation	Initiates therapy topics, engages in co-regulation, takes ownership of healing journey
5. Integrated Participation	Full presence in life: emotionally, socially, spiritually, physically. Able to navigate stress while staying regulated.	Flexible Autonomic Regulation	Adapts to stress, shows resilience, participates in meaningful occupation, rebuilds identity

Clinical Insight

- Clients may fluctuate between levels depending on stress, safety, or triggers.
- Participation is not linear – regression is expected and normal.
- Assessing the client's engagement level is critical before choosing tasks or therapeutic goals.

Therapeutic strategy per level

Level	Therapist Focus
Withdrawal	Safety, co-regulation, basic sensory grounding
Passive	Rhythm, relational cues, bottom-up presence
Task	Structure, success experiences, pacing
Relational	Emotional exploration, identity, social repair
Integrated	Meaning-making, self-agency, occupation reintegration

5-Stage Model of Physical Engagement

A nervous system-based participation ladder

Level	Client State	Nervous System Profile
1. Protective Inhibition	Body is in full defense mode. Movement is perceived as threat. Pain dominates. Client avoids or resists all physical input.	Dorsal vagal + sympathetic freeze Safety cues, sensory grounding, relational trust
2. Tolerated Passivity	Client allows touch or movement, but remains passive. Not yet initiating or trusting the process. Still externally driven.	Mixed autonomic state Gentle activation, rhythm, education on pain science
3. Conditioned Activation	Begins to move or follow exercise protocols with effort. Progress is goal-oriented but still fear-based. Motivation often linked to avoiding setbacks.	Sympathetic + fragile ventral Build tolerance, strengthen control, reframe pain
4. Responsive Engagement	Body and mind begin to trust movement. Initiates, adjusts, and reflects on experience. Safety is returning.	Ventral vagal activation Encourage autonomy, explore movement variety
5. Embodied Resilience	Movement is integrated. Client adapts with awareness. Injury or pain no longer dictates identity.	Flexible autonomic regulation Expand capacity, reintegrate into life roles or sport

Relevant Theories This Aligns With:

- Graded Exposure in Chronic Pain (Butler & Moseley)
- Biopsychosocial Model
- Pacing & Graded Activity in energy management
- Trauma-informed physio approaches (e.g. Carolyn Vandyken, Neil Pearson)
- Polyvagal-informed rehab frameworks

Clinical Tip:

If movement feels like threat, the client won't perform it – no matter how safe it is biomechanically.

- Therefore, nervous system readiness is a precondition for progress in physiotherapy.

5-Stage Psychological Participation Model

A polyvagal-informed map of therapeutic readiness

Stage	Client Experience	Nervous System Profile	Therapy Presentation
1. Survival Mode / Disengaged	Client appears shut down, disconnected, or ambivalent. Words are minimal or guarded. Insight is absent or intellectualized.	Dorsal vagal shutdown or sympathetic overwhelm	"I don't know." "I'm tired." Silence, monotone, minimal eye contact
2. Compliant but Defended	Attends sessions, but emotional engagement is limited. Uses defense mechanisms (e.g. humor, avoidance, perfectionism).	Mixed state: low ventral, high sympathetic	Over-thinks, avoids emotions, appears "fine" but stays surface-level
3. Emotionally Reactive / Testing Boundaries	Begins showing raw emotion, resistance, or frustration. Emotions emerge without regulation. Therapist seen as both threat and lifeline.	Sympathetic fight/flight + fragile ventral	"Why do I feel worse?" Anger, tears, splitting, dependency or detachment
4. Co-Regulated Curiosity	Engages in reflective dialogue, explores inner states. Begins to understand emotional patterns. Willing to feel discomfort.	Ventral vagal activation	"I noticed this about myself." Can stay with emotion, tolerate repair, own insight
5. Self-Directed Integration	Engages in reflective dialogue, explores inner states. Begins to understand emotional patterns. Willing to feel discomfort. Acts on insight outside of sessions. Regulates emotions independently. Seeks growth over symptom control. Trauma has meaning, not just pain	Flexible autonomic regulation	Can stay with emotion, tolerate repair, own insight Brings material to sessions, sets goals, initiates rupture repair, embodies change

Therapist Goals Per Stage:

Stage Focus

1. Survival – Provide co-regulation, build safety, attune nonverbally
2. Compliant – Invite authenticity, name defenses gently, avoid premature interpretation
3. Reactive – Hold emotion safely, validate, model regulation
4. Curious – Deepen reflection, support meaning-making, offer cognitive tools
5. Integrated – Encourage embodiment, expand identity, support long-term vision

Aligns With:

- Polyvagal Theory in Therapy (Deb Dana)
- Stages of Change (Prochaska & DiClemente)
- Window of Tolerance (Dan Siegel)
- IFS / Parts Work (Richard Schwartz)
- Attachment Theory – especially as relational safety is built over time

Clinical Insight:

The mind opens when the body feels safe. Insight without safety creates disintegration, not healing.

Part 6

Tools to Track The ANS and Regulation

Why Assess Before Treating?

"You cannot regulate what you do not recognize."

Why Assess Before Treating?

In trauma-informed nervous system work, assessment is not about diagnosis — it's about mapping patterns. We ask:

- What state is dominant?
- What triggers state shifts?
- What's the client's baseline rhythm?
- Where is the entry point for safety?

Assessment should reveal how the body organizes around survival, and what regulation strategy will meet the system where it is, not where we want it to be.

Core Principles of Nervous System Assessment

- States over symptoms – Look at physiological patterns, not just mood or behavior
- Function over form – What is this dysregulation doing for the system? What's it protecting?
- Capacity over compliance – Can the client shift states, or are they stuck?
- Regulation before cognition – Don't over-analyze what hasn't yet stabilized

Key Tools You'll Learn to Use

Tool	What It Screens	Application
State Tracking Worksheet	Dominant state (sympathetic, dorsal, ventral)	Daily/weekly state awareness
Breath Pattern Checklist	Dysregulated breath types (freeze, panic, over-breathing)	Match breath to state and intervention
HRV & BOLT Tracking Sheet	Nervous system trends over time	Use with wearables or manual breath test
Interoceptive Awareness Scale	Capacity to feel inside the body	Predicts how a client will respond to bottom-up tools
Therapist Observational Markers & Window of tolerance	Tone, posture, eye contact, voice, movement	Session-based nervous system cues

These tools form a clinical mirror — showing where to begin, what to avoid, and when a client is ready for deeper work.

Integration with the Protocol (Coming in Module 2)

These assessments are not separate from intervention — they're part of the flow:

1. Observe the state
2. Identify the entry point (breath, movement, eyes, etc.)
3. Apply rhythmic regulation
4. Reassess (before > after comparison)
5. Build tolerance and coherence over time

Assessment is not a one-time event — it's a rhythm of listening, noticing, and adjusting.

References

- Dana D. Polyvagal Theory in Therapy. W.W. Norton; 2018.
- Porges SW. Clinical Applications of the Polyvagal Theory. W.W. Norton; 2014.

HRV tracking

A “snap-shot” of the ANS and vagal tone.

Tools to Track Regulation: HRV Devices, BOLT Score, and Clinical Observation

“What we can measure, we can map. What we can map, we can shift.”

Why Tracking Matters

Nervous system regulation is often invisible until it's not. By tracking physiological markers, we move:


- From guessing to grounding
- From symptoms to states
- From talk about behavior to insight about rhythm

These tools help make the invisible visible — for both the therapist and the client.

Heart Rate Variability (HRV) Device

<https://www.heartmath.com/neurosync>

Use this link if you are interested in purchasing Heartmath emwave for your practice



**30+ YEARS OF PIONEERING RESEARCH
IN HEART-BRAIN SCIENCE**

BOLT - Body Oxygen Level Test


A functional window into CO₂ tolerance, vagal tone and interoception.

The BOLT Score (Body Oxygen Level Test)

The BOLT score is a low-tech, highly effective breath hold test that reflects:

- CO₂ tolerance
- Breathing efficiency
- Autonomic nervous system flexibility

It helps identify hidden hyperventilation, sympathetic overdrive, and low vagal tone — all common in chronic stress.



BOLT Score (Seconds)	Interpretation
< 10 sec	High dysregulation / over-breathing
10-20 sec	Moderate imbalance
20-30 sec	Near functional balance
> 30 sec	Good breath and vagal regulation

Higher BOLT = more resilience, safer access to parasympathetic tone, and better baseline coherence.

BOLT Score Interpretation Table

BOLT Score (sec)	Nervous System Interpretation	Regulation Strategy
<10 sec	High sympathetic tone, likely overbreathing	Begin with short breath holds + nasal breathing only
10–20 sec	Moderate stress load, early fatigue	Coherence breathing, light aerobic movement
20–30 sec	Functional baseline	Build into nose breathing habits, light cardio
30+ sec	Excellent CO ₂ tolerance	Body is adaptable, resilient — maintain protocol

BOLT score is an indirect reflection of chemosensitivity and vagal tone.

Clients with low scores often have poor HRV, shallow breathing, and a narrow window of tolerance. Improving BOLT = improving nervous system resilience.

HRV & BOLT Tracking

HRV & BOLT Tracking

“Track the rhythm, don’t chase the number.”

Why Track HRV and BOLT?

These two markers — Heart Rate Variability (HRV) and Body Oxygen Level Test (BOLT) — give objective insight into the nervous system’s adaptability. When tracked over time, they help clinicians and clients:

- See patterns of recovery or regression
- Monitor progress through regulation work
- Identify early signs of overload or shutdown
- Match protocols more precisely to physiological need

Tracking creates a mirror for rhythm and resilience.

Tip: Use the same device each time for consistency. Use trends to guide, not single readings.

Optional: Add “Coherence Level” if using emWave or Inner Balance.

BOLT Interpretation Guide

BOLT Score

Meaning

< 10 sec

Likely over-breathing / stress overload

10–20 sec

Moderate dysfunction / transition zone

20–30 sec

Functional, improving regulation

> 30 sec

Strong vagal tone, high CO₂ tolerance

Therapist Tips for Use

- Use HRV/BOLT pre- and post-intervention (breathwork, eye movement, somatic work)
- Match progress to subjective state change for integrated insight
- Celebrate small wins (e.g. from 8 to 12 sec BOLT = big nervous system shift)
- Adjust protocols based on plateaus or regressions
- Teach clients to self-track once they show interest or ownership

References

- McCraty R, Zayas M. Heart rhythm coherence and self-regulation. *Front Psychol.* 2014;5:1090.
- Buteyko KP. *The Buteyko Method: Clinical Application.* 1991.
- Lehrer P, Gevirtz R. HRV biofeedback: How and why it works. *Front Psychol.* 2014;5:756.

HRV & BOLT Symptom Checklist: Clinical Signs of Nervous System Dysregulation

“When coherence is low, the body whispers in symptoms.”

Why This Matters

Not all clients can access biofeedback tools. But trained clinicians can read HRV and BOLT patterns by observing symptoms, breath behavior, and regulation capacity.

This checklist helps therapists match subjective symptoms to likely physiological rhythms, especially in the absence of HRV tracking devices.

HRV Symptom Patterns

HRV Pattern	Likely Symptom Clusters	Interpretation
Low HRV	- Fatigue- Low motivation- Poor stress recovery- Emotional flatness- GI sluggishness- Chronic pain or burnout	Suggests vagal suppression, possible dorsal vagal dominance, or allostatic overload
Erratic HRV	- Emotional swings- Panic episodes- Sensory hypersensitivity- Difficulty focusing- Autonomic instability	Often seen in trauma survivors or ADHD clients; signals poor ANS rhythm and oscillation
High but incoherent HRV	- Overbreathing- Compulsive calmness- Disconnected from body- “Functional freeze”	May reflect sympathetic control masking deeper dysregulation
High + Coherent HRV	- Grounded energy- Calm alertness- Resilient recovery- Vagal flexibility	Ideal state; good predictor of therapeutic receptivity and long-term resilience

BOLT Score Symptom Correlation

BOLT Score Range	Common Signs	Nervous System Insight
< 10 sec	- Mouth breathing- - Sleep dysfunction- - Morning headaches- - Low patience- - Emotional reactivity	Indicates hyperventilation, sympathetic drive, poor CO ₂ tolerance
10-20 sec	- Variable sleep- - Improved focus- - Better energy with rest- - Reactivity under pressure	Moderate regulation with room for growth
20-30 sec	- Calm focus- - Good energy- - Consistent recovery- - Emotional resilience	Functional breathing and vagal balance
> 30 sec	- Stable moods- - Restful sleep- - Clear mind under stress- - Body-mind coherence	Reflects strong parasympathetic tone and low baseline arousal

References

- McKeown P. The Oxygen Advantage. HarperCollins; 2016.
- Shaffer F, Ginsberg JP. An overview of HRV metrics and norms. Front Public Health. 2017;5:258.
- Porges SW. Polyvagal Theory. W.W. Norton; 2011.
- HeartMath Institute. Science of the Heart, Vol. 2. HeartMath Institute; 2016.
- Buteyko KP. Breathing for Children. Buteyko Clinic; 1993.
- McCraty R. Science of the Heart, Vol. 2. HeartMath Institute; 2016.
- Lehrer PM, Vaschillo E, Vaschillo B. Heart rate variability biofeedback: methodology and outcomes in children. Appl Psychophysiol Biofeedback. 2010;35(3):193-198.
- McCraty R, Childre D. Heart-brain communication. Integr Med Clin J. 2010;9(3):20-26.
- Lehrer PM, Gevirtz R. HRV biofeedback and psychophysiological resilience. Appl Psychophysiol Biofeedback. 2014;5:756.

HRV & BOLT Scores in Children: Interpreting Regulation in Growing Systems

Why Children Need Different Metrics

Children are not just smaller adults. Their autonomic nervous systems:

- Are still under construction
- Rely more on co-regulation
- Reflect environmental safety more acutely
- Show wider variability in HRV and breath patterns

Tracking regulation in children requires softer markers, developmental insight, and the ability to read play, posture, and voice tone as part of the picture.

HRV in Children

Baseline HRV tends to be higher in younger children due to faster heart rates and increased vagal oscillations. However:

- Erratic HRV may reflect overstimulation or attachment trauma
- Flat or suppressed HRV may reflect chronic stress, shutdown, or sensory overload
- High coherence in a calm emotional state is ideal, but less common in early childhood

BOLT Score in Children

The BOLT score reflects breath-hold comfort after a normal exhale. In children, it:

- Indicates CO₂ sensitivity
- Shows vagal-brainstem resilience
- Is influenced by mouth breathing, asthma, anxiety, or early trauma

Signs of low BOLT:

- Mouth breathing
- Poor sleep quality
- Frequent sighing
- Short fuse / emotional flooding
- ADHD-like symptoms (linked to over-breathing)

"A child who can't hold their breath often can't hold their state."

References

- McKeown P. The Oxygen Advantage. HarperCollins; 2016.
- Shaffer F, Ginsberg JP. An overview of HRV metrics and norms. Front Public Health. 2017;5:258.
- Porges SW. Polyvagal Theory. W.W. Norton; 2011.
- HeartMath Institute. Science of the Heart, Vol. 2. HeartMath Institute; 2016.
- Buteyko KP. Breathing for Children. Buteyko Clinic; 1993.
- McCraty R. Science of the Heart, Vol. 2. HeartMath Institute; 2016.
- Lehrer PM, Vaschillo E, Vaschillo B. Heart rate variability biofeedback: methodology and outcomes in children. Appl Psychophysiol Biofeedback. 2010;35(3):193-198.

Interoception

Clinical Observation & Interoception

Sometimes the best tracking tool is your eyes and ears.

Observation	Possible Interpretation
Shallow breathing, mouth open	Sympathetic overactivation, low CO ₂ tolerance
Sighing, breath-holding	Dorsal shutdown or unresolved tension
Monotone voice, flat affect	Low vagal tone, freeze
Excessive blinking / darting eyes	Hypervigilance, sympathetic state
Hand-wringing, leg bouncing	Incomplete mobilization, fawn state
Teariness during grounding	State transition, thawing pattern
Sudden stillness + staring	Shutdown response or freeze trigger

References

- McKeown P. The Oxygen Advantage. HarperCollins; 2016.
- Shaffer F, Ginsberg JP. An overview of HRV metrics and norms. Front Public Health. 2017;5:258.
- Porges SW. Polyvagal Theory. W.W. Norton; 2011.
- HeartMath Institute. Science of the Heart, Vol. 2. HeartMath Institute; 2016.

Interoceptive Awareness Scale: Tracking the Body's Inner Language

“Interoception is the doorway to self-regulation.”

What Is Interoception?

Interoception is the ability to sense the internal state of the body — heartbeat, breath, hunger, temperature, tension, and emotion-in-body.

Healthy interoception allows clients to:

- Detect when they're stressed or overwhelmed
- Notice tension before it becomes pain
- Feel emotional shifts before they escalate
- Respond with regulation tools in real-time

In trauma, interoception is often disrupted — either:

- Blunted (numb, disconnected)
- Hyperactive (overwhelmed, flooded)

References

1. Mehling WE et al. The Multidimensional Assessment of Interoceptive Awareness (MAIA). PLoS One. 2012;7(11):e48230.
2. Craig AD. How do you feel—now? The anterior insula and human awareness. Nat Rev Neurosci. 2009;10(1):59-70.
3. Badenhorst C. NeuroSync Interoceptive Training Notes. 2022-2025.

Interoceptive Awareness Self-Assessment

Rate each of the following on a scale of **0 to 3**:

Item	Description	Score (0-3)
1. I notice my heartbeat without touching my chest		
2. I can tell when I'm holding my breath		
3. I feel my stomach tighten when stressed		
4. I can detect changes in my body before I feel anxious or upset		
5. I notice when my breathing changes with emotion		
6. I feel connected to my body during the day		
7. I recognize when I'm hungry or full without checking the time		
8. I can tell when I need rest, even before I feel tired		
9. I feel emotions as physical sensations in my body		
10. I use body awareness to help myself calm down		

Scoring Key:

- 0 = Not true at all
- 1 = Rarely true
- 2 = Sometimes true
- 3 = Often true

Interpretation

Total Score Range	Interoceptive Profile
0-10	Blunted interoception: Likely disconnected from body, needs slow bottom-up work
11-20	Developing awareness: Some capacity to feel inside, regulation tools may begin
21-30	Moderate awareness: Ready to use body as a guide in therapy
31-40	High interoception: Ideal for deeper somatic, breath, and emotional integration work

Breath pattern tracking

Breath Pattern Checklist

“Breath is the story the body tells before the mind catches up.”

Why Breath Matters in Assessment

Breathing is the most immediate expression of nervous system state — faster than words, often outside awareness, and deeply tied to trauma patterns.

By observing breath, we can:

- Detect survival responses
- Predict regulation capacity
- Match the right intervention
- Support state transitions

This checklist offers a clinical way to read the breath and identify which regulation strategies are most appropriate.

Breath Pattern Identification Table

Interpretation Tips

- A collapsed breath pattern may indicate shutdown even if the person appears calm
- Fast or loud breathing often signals mobilization or trauma activation
- Mouth breathing should be addressed early; it limits vagal tone and distorts HRV
- Pattern + state awareness guides the choice of breathwork (see Module 2)

Therapist Use

- Observe the client's breath at session start, during stress talk, and after grounding
- Track how breath shifts with emotional activation or interventions
- Reflect: “Is this breath a sign of defense or release?”

Breath Pattern	Description	Common in This State	Clinical Notes
Shallow chest breathing	Fast, high in chest	Sympathetic, anxiety	Often accompanied by rapid speech, clenched jaw
Breath holding	Long pauses, no exhale	Freeze, shutdown	Common in trauma survivors, even during sleep
Mouth breathing	Constant oral inhalation	Stress, poor CO ₂ tolerance	Linked to low BOLT, sleep dysfunction, brain fog
Sighing or gasping	Frequent sighs or yawns	Hidden hyperventilation	Body trying to reset CO ₂ – may reflect dorsal/overload
Over-breathing	Deep but too frequent	High sympathetic arousal	Not always calming; can increase anxiety
Silent stillness	No visible movement	Freeze or dissociation	Often misread as “calm” – observe context
Long exhale with tone	Regulated vagal rhythm	Ventral vagal state	Sign of safety and autonomic flexibility

References

1. McKeown P. The Oxygen Advantage. HarperCollins, 2016.
2. Jerath R et al. The physiological basis of breathing techniques. Int J Yoga. 2006;1(1):2-7.
3. Porges SW. The Pocket Guide to the Polyvagal Theory. Norton, 2017.
4. Badenhorst C. NeuroSync Clinical Observations. 2021-2025.

Example Clinical Matches

Client Presentation	Likely Breath Pattern	Suggested Focus
Panic attacks	Over-breathing, gasping	Restore CO ₂ tolerance, extended exhale
Chronic fatigue	Mouth breathing, silent stillness	Nasal re-training, rhythmic humming
Dissociation	Breath-holding, no rhythm	Gentle rocking with paced nasal breath
ADHD/irritability	Shallow chest breath	Slower diaphragm breathing, coherence work

Window of tolerance tracking

Window of Tolerance Tracking Scale: Mapping Your Nervous System in Real Time

“The window is not fixed — it widens with rhythm, breath, and safety.”

What Is the Window of Tolerance?

Coined by Dan Siegel, the Window of Tolerance is the range of emotional and physiological arousal within which a person can function optimally — think clearly, connect socially, and respond flexibly.

When outside this window, individuals move into:

- Hyperarousal (Sympathetic) – Anxiety, panic, agitation
- Hypoarousal (Dorsal Vagal) – Numbness, shut down, disconnection

This scale helps clients track their daily state, notice patterns, and build awareness of what shrinks or widens their window

Window of Tolerance Self-Rating Scale

Instructions:

Circle or shade the zone that best reflects how you felt during each time period. Use one row per day.

Date	Morning	Midday	Evening	Main Trigger	Main Anchor	Notes
						
						

State Key:

- **Freeze / Shut Down** – Numb, foggy, disconnected
- **In the Window (Optimal Zone)** – Present, grounded, focused
- **Hyperarousal** – Anxious, racing, flooded
- **Overwhelm / Collapse** – Exhausted, impulsive, dissociating

Clinical Use

Observation Area	In the Window	Hyperaroused	Hypoaroused
Thinking	Clear, flexible	Racing, scattered	Blank, foggy
Emotions	Balanced, flowing	Flooded, reactive	Numb, unreachable
Body	Grounded, steady	Tense, jittery	Limp, disconnected
Breathing	Slow, rhythmic	Fast, shallow	Held, minimal
Voice	Expressive, stable	Loud, fast	Quiet, flat

Use this during:

- Intake sessions
- Pre/post intervention tracking
- Client reflection journals
- Group therapy to normalize shifts

Window of Tolerance Tracking Scale: Mapping Your Nervous System in Real Time


Every emotion you experience has an effect on your body and on your resilience, whether or not you are aware of it. Every emotion immediately causes changes in your body, affects your ability to build and sustain your energy and either renews or depletes your resilience. The two main physiological systems that control the cascade of these changes in your body are the hormonal system and the autonomic nervous system (ANS).

Scale	Description	Sensations	Possible Interventions
10	High anxiety or panic	Racing heart, jittery, rapid breath	Breath pacing, orienting, cold water
9	High alert	Over-focused, scanning for threat	5-4-3-2-1 grounding, slow exhale
8	Irritable/stressed	Jaw tension, inner pressure	Shake it out, movement, sighs
7	Activated but functional	Busy mind, energised	Coherence breath, posture reset
6	Alert and aware	Mild effort, focused	Flow state zone – support
5	Grounded and calm	Clear, steady, connected	Maintain – reflect, rest, act
4	Soft focus	Relaxed, open, creative	Great for healing, journaling
3	Drowsy or spaced	Wandering, yawning	Gentle alerting – stretch
2	Disconnected	Numb, low sensation	Touch object, orient with eyes
1	Frozen or collapsed	Can't move, hopeless	Warmth, movement, safe cue
0	Shutdown	Gone blank, can't engage	Co-regulate, don't push – stay close

Zone	WOT Status	Tracking Scale Range	Nervous System State
■ Optimal Zone	Within WOT	4–6	Ventral vagal calm, engagement, flow
▲ Hyperarousal	Above WOT	7–10	Sympathetic fight-flight activation
▼ Hypoarousal	Below WOT	0–3	Dorsal vagal freeze/shutdown


Purpose of the Scale:

1. Transforms a vague concept (WOT) into a usable daily self-check
2. → Clients often say “I don’t know where I am.” The scale gives them language: “I’m at a 2 – shut down” or “I’m at an 8 – overwhelmed.”
3. Empowers nervous system literacy
4. → Clients begin to notice: “When I hit 3, I stop responding. That’s my red flag.”
5. Tracks state shifts in real time
6. → Therapists can ask before/after breathwork: “Where were you before? Where are you now?”
7. Links state to intervention
8. → Each zone includes suggested strategies, so the scale becomes a built-in guide to regulation.



References

1. McCraty R, Zayas MA. Emotional State Mapping via HRV. HeartMath Institute, 2014.
2. Badenhorst C. NeuroSync Regulation Matrix. 2023-2025.
3. Porges SW. Clinical Applications of the Polyvagal Theory. Norton, 2014.



Every emotion you experience has an effect on your body and on your resilience, *whether or not you are aware of it*. Every emotion immediately causes changes in your body, affects your ability to build and sustain your energy and either renews or depletes your resilience. The two main physiological systems that control the cascade of these changes in your body are the hormonal system and the autonomic nervous system (ANS).

Part 7

Module 2

Practical Nervous System Regulation Techniques

Practical Nervous System Regulation Techniques

Learning Objectives & Outcomes

By the end of this module, participants will be able to:

1. Explain the scientific mechanisms behind bottom-up nervous system regulation, including breath, eye movements, somatic input, and sensory integration.
2. Identify the most appropriate regulation techniques for different nervous system states (e.g. sympathetic hyperarousal, dorsal hypoarousal, freeze/fawn cycles).
3. Match specific techniques to clinical presentations such as anxiety, PTSD, ADHD, IBS, chronic pain, and emotional dysregulation.
4. Demonstrate accurate use of selected breathwork, eye movement, and somatic tools in real or simulated clinical settings.
5. Recognize client readiness and contraindications for each technique, using observation-based assessment.
6. Sequence and layer regulation interventions based on a client's nervous system profile and therapeutic goals.
7. Build a full-session regulation protocol using the NeuroSync Regulation Flow Framework.
8. Evaluate the effect of interventions using symptom tracking, HRV/BOLT data, and client-reported outcome measures.

Expected Competencies by Completion

Participants will be able to:

- Use at least three different breathing strategies, adjusted to vagal tone and clinical state.
- Apply two or more eye movement techniques, including vertical, lateral, and gaze anchoring, for emotional looping and cortical regulation.
- Guide clients through basic somatic grounding and proprioceptive engagement.
- Identify at least five signs of dysregulation or readiness that influence when to initiate or pause techniques.
- Construct a layered protocol for at least three clinical case types (e.g. trauma client, ADHD child, chronic pain adult).
- Demonstrate familiarity with the BOLT test, HRV tracking, and state mapping for feedback-informed treatment.
- Reflect critically on ethical and safety considerations in bottom-up interventions.

Why Regulation Must Be Felt, Not Forced

"Regulation is not a toolset — it is a rhythm we teach the body to remember."

"You cannot talk a nervous system into safety. You have to show it."

The Bridge from Awareness to Action

Module 1 gave you maps — states, windows, loops, and signals.

Now, in Module 2, you begin guiding the system.

But nervous system regulation is not a cognitive task.

It's not something we convince the body to do — it's something we help it remember.

This is the core of bottom-up work:

We don't just "calm the mind."

We create conditions for the body to access safety — and from there, the mind follows.

Why Top-Down Alone Doesn't Work

Trauma, chronic stress, and illness often disconnect the body from awareness.

In these states:

- The prefrontal cortex is offline
- Coherence is low
- The vagus nerve is suppressed
- Logical thinking feels unsafe or inaccessible

This is why traditional interventions (e.g. CBT, insight, planning, advice) often don't work for:

- Anxiety that overrides breathing
- Trauma loops that replay without reason
- Children who "freeze" during schoolwork
- Adults with shutdown or chronic pain

Bottom-Up = Regulation First

- Bottom-up tools — such as breath, eye movement, grounding, rhythm, and body-based awareness — bypass cortical thinking and access brainstem and limbic regulation first.

This allows:

- A return to the Window of Tolerance
- Reconnection to social engagement systems
- Activation of vagal tone and interoception
- Re-entry into executive functioning (thinking, planning, reflecting)

Your Role as a Therapist

You're not here to fix dysregulation.

You're here to help the system reclaim rhythm.

This means:

- Slowing down
- Observing signs of readiness
- Guiding simple inputs: breath, gaze, posture, rhythm
- Letting the body speak first — before asking the mind to respond

Breath Science & Nervous System Regulation

"Breathing is the only autonomic function we can consciously control — and through it, we regulate every other system."

The Power of Breath in Regulation

Breath is the most direct and immediate intervention for shifting nervous system states. It engages both voluntary and autonomic pathways, influencing:

- Heart rate variability (HRV)
- Vagal tone
- Blood gases (CO₂, O₂)
- Emotional state
- Brainwave patterns
- Immune and inflammatory responses

Unlike other autonomic functions, breathing can be controlled intentionally, allowing us to downregulate threat responses and access the ventral vagal state of safety.

Physiological Basis of Breath Regulation

1. Autonomic Nervous System

- Inhalation stimulates the sympathetic branch, increasing alertness and heart rate.
- Exhalation activates the parasympathetic branch, especially via the vagus nerve, slowing the heart rate and supporting rest and repair (1).

2. Vagal Afferents

Slow, diaphragmatic breathing activates vagal afferent fibers in the lungs and diaphragm, sending safety signals to the nucleus tractus solitarius (NTS) in the brainstem, which integrates with the insula, amygdala, and prefrontal cortex (2,3).

3. Respiratory Sinus Arrhythmia (RSA)

RSA is the natural variation in heart rate during the breathing cycle:

- Heart rate increases on inhalation
- Heart rate decreases on exhalation
- This phenomenon is a marker of vagal tone and emotional resilience (4).

4. Gas Exchange and CO₂ Sensitivity

CO₂ is not merely a waste gas — it plays a crucial role in chemoreceptor sensitivity, anxiety modulation, and cerebral blood flow. Low CO₂ tolerance can exacerbate anxiety, fatigue, and cognitive shutdown (5). Breathwork improves CO₂ tolerance and resets the body's chemical sensitivity to safety.

Breath and Brain Function

Breath impacts brain oscillations and cortical states:

- Fast, shallow breathing correlates with beta waves → alertness, anxiety
- Slow, rhythmic breathing promotes alpha and theta waves → calm focus, memory consolidation (6)
- Breath synchronizes activity in the prefrontal cortex, limbic system, and brainstem, enhancing self-regulation and emotional control (7)

Multi-System Integration

System	Breath-Related Regulation
Cardiovascular	Heart rate, HRV, blood pressure via baroreceptor stretch
Endocrine	Cortisol reduction, melatonin production through vagal tone (8)
Immune	Anti-inflammatory cytokines via vagal activation (9)
Digestive	Gut motility and enzyme release through parasympathetic dominance
Cognitive	Enhanced executive function via improved prefrontal access (10)

Clinical Insight:

For clients in sympathetic overdrive, dorsal shutdown, or trauma loops, conscious breathwork becomes a portal to safety and agency:

- Breath is non-invasive, portable, and immediate
- Breath can be taught, tracked, and refined across sessions
- Breath often reveals dysregulation before the client has words for it

"When you change the breath, you change the chemistry. When you change the chemistry, you change the story."

References

1. Jerath R, Edry JW, Barnes VA, Jerath V. Physiology of long pranayamic breathing: Neural respiratory elements may provide a mechanism that explains how slow deep breathing shifts the autonomic nervous system. *Med Hypotheses*. 2006;67(3):566-71.
2. Porges SW. The polyvagal theory: Neurophysiological foundations of emotions, attachment, communication, and self-regulation. W. W. Norton & Company; 2011.
3. Critchley HD, Harrison NA. Visceral influences on brain and behavior. *Neuron*. 2013;77(4):624-38.
4. Lehrer PM, Gevirtz R. Heart rate variability biofeedback: how and why does it work? *Front Psychol*. 2014;5:756.
5. Courtney R. The functions of breathing and its dysfunctions and their relationship to breathing therapy. *Int J Osteopath Med*. 2009;12(3):78-85.
6. Zaccaro A, et al. How breath-control can change your life: A systematic review on psycho-physiological correlates of slow breathing. *Front Hum Neurosci*. 2018;12:353.
7. Ma X, Yue ZQ, Gong ZQ, et al. The effect of diaphragmatic breathing on attention, negative affect and stress in healthy adults. *Front Psychol*. 2017;8:874.
8. Brown RP, Gerbarg PL. Yoga breathing, meditation, and longevity. *Ann N Y Acad Sci*. 2009;1172:54-62.
9. Tracey KJ. The inflammatory reflex. *Nature*. 2002;420(6917):853-9.
10. Thayer JF, Lane RD. Claude Bernard and the heart-brain connection: Further elaboration of a model of neurovisceral integration. *Neurosci Biobehav Rev*. 2009;33(2):81-8.

Breath

Breath is the only system that bridges body, brain, and emotion — in real time...

Breath is like medicine – not all created for the same purpose

Reflection: Breath as Soul Language

Breath brings you out of hiding.

Breath gives you the courage to be seen again.

Breath is an unseen language — the most powerful feeder of body and soul. It is both mechanical and chemical, yet deeply sacred. It is pure soul food.

Each inhale and exhale regulates coherence in real time. Breath is the very molecule of the unseen, the bridge between spirit and cell. It carries the key to open and close the body's gates — shifting the autonomic nervous system between fight, flight, freeze, and safety.

Breath is like a key to a door. We breathe to center ourselves before we engage. With each inhale and exhale, breath creates a brooding space — a quiet atmosphere where restoration can emerge, ideas can take root, healing can begin, and identity can safely unfold.

I see it like a hen sitting on her eggs. She doesn't force life to happen — she simply creates the right environment for life to grow. Breath does the same for us: it incubates possibility.

Breath can clear a stuck mind, unlock rational thought, and shift gears from amygdala hijack or cortical chaos into balance, freedom, identity, and self-control — the kind of self-control that lets your authentic self move from survival to creation.

Trauma often closes this doorway. It makes us hold back from the world, brace, and shrink into sympathetic overdrive. But breath — in its gentle flow and unwavering rhythm — still holds the capacity for breakthrough.

Even at a cellular level, breath fuels life. Our mitochondria thrive on it, using oxygen to create the very energy that keeps us alive (85% of our ATP — the body's energy currency — comes from oxygen).

In nervous system regulation and therapy, oxygen delivery isn't just biochemistry. It's foundational.

Every breath you take is a chance to open the door again.

Breath Science & Regulation: Why Breathing Is the Gateway

Why Breath Comes First

Breath is the only autonomic function that we can consciously control – and by doing so, we gain a direct handle on:

- Heart rate
- Vagal tone
- Blood pH and CO₂ levels
- Emotional arousal
- Cortical access

It is the fastest, most accessible way to shift nervous system state, even in dysregulation.

The Science of Breath & the Nervous System

System	Breath Interaction
Autonomic Nervous System (ANS)	Inhalation activates sympathetic (arousal), exhalation activates parasympathetic (calm).
Vagus Nerve	Slow, nasal, diaphragmatic exhalations stimulate vagal afferents to the brainstem.
Cardiac Function	Breath influences Heart Rate Variability (HRV) via Respiratory Sinus Arrhythmia (RSA).
Brainstem & Cortical Balance	Rhythmic breathing synchronizes brain oscillations and improves executive function.
Chemistry & CO ₂	Breath regulates blood gases – especially CO ₂ , which affects anxiety, alertness, and fatigue.

Respiratory Sinus Arrhythmia (RSA)

RSA is the natural increase in heart rate during inhalation and decrease during exhalation.

This variation — when coherent — reflects optimal vagal function and emotional regulation.

- Inhale = slight sympathetic activation
- Exhale = parasympathetic rebound
- Longer, smoother exhalations = greater vagal tone

This is the physiological signature of safety.

Breath & Brain States

Breathing affects brainwave activity:

- Fast, shallow breathing → Beta → Anxiety, alertness
- Slow, nasal breathing → Alpha/Theta → Calm focus, introspection
- Extended breath holds → Theta → Dissociation or deep trance (caution in trauma clients)

Breath = Chemical, Mechanical, and Emotional

Breathing isn't just about oxygen — it changes:

- CO₂ levels → tolerance affects panic/fatigue
- Baroreceptor stretch → regulates blood pressure and vagal input
- Muscle and diaphragm feedback → activates sensory-motor brain areas
- Emotional encoding → breath patterns link directly to trauma and memory

"The way we breathe is the way we feel. And the way we feel is the way we live."

Baro = pressure

Reflex = automatic response

The baroreflex is your body's built-in pressure sensor that helps keep your heart, blood pressure, and nervous system stable. When we breathe slowly, we 'massage' these sensors, which tells your brain that you're safe — allowing your whole system to relax."

Types of Breathing Techniques: What to Use, When and Why

“Not all breathing calms. Some breath awakens, some centers, and some clears trauma.”

Extended Exhalation Breathing

“The body exhales what the mind cannot. In the lengthening of the breath, safety returns.”

Scientific Basis

Extended exhalation breathing activates the parasympathetic nervous system, specifically through the vagal afferents in the lungs, diaphragm, and thoracic cavity. During a prolonged exhale, baroreceptors in the heart and lungs send signals to the nucleus tractus solitarius (NTS) in the brainstem, modulating activity in the amygdala, insula, and prefrontal cortex (1,2). It also increases Respiratory Sinus Arrhythmia (RSA) — a key marker of vagal tone and emotional flexibility — by deepening the heart rate variability associated with the breath cycle (3). The longer the exhale, the stronger the vagal signal.

Clinical Purpose

- Reduce sympathetic overactivation
- Down-regulate fight/flight loops
- Shift out of hyperarousal or panic
- Reset heart-brain synchronization
- Access the ventral vagal state
- Support emotional processing without overwhelm

This technique is foundational for clients who are “stuck on ON” — hyper-alert, anxious, agitated, or unable to rest.

How to Do It

Simple pattern:

Inhale gently through the nose for 4 seconds

Exhale slowly and completely through the nose or mouth for 6–8 seconds

Optional: brief 2–3 second pause at the bottom of the exhale

Key coaching cues:

- “Breathe into the belly, soft and low.”
- “Let the exhale take a little longer than the inhale.”
- “Imagine your breath melting down your spine as you exhale.”

Advanced variation:

Add gentle pursed lips on the exhale to create slight resistance, which increases baroreceptor activation and vagal tone (4).

Observable Signs of Regulation

- Shoulders drop, face softens
- Eye contact returns
- Decrease in verbal speed or tone
- Hands unclench, posture releases
- HRV begins to increase (if measured)
- Client reports: "I feel a bit more here," "It's slowing down," or "I can think again"

Best Matches Per Diagnosis / Symptom Cluster

Diagnosis / Pattern	Suitability
Generalized anxiety disorder	✓ Excellent
Panic disorder	✓ With caution (begin short)
PTSD (hyperarousal type)	✓ Foundational
ADHD (high sympathetic)	✓ As prep tool
Dissociation / dorsal vagal freeze	✗ Use with caution; may increase shut down
Insomnia, burnout, stress overload	✓ Core technique
Autism (with hyperreactivity)	✓ Introduce gradually with sensory support

References

1. Porges SW. The Polyvagal Theory. W. W. Norton; 2011.
2. Critchley HD, Garfinkel SN. Interoception and emotion. Curr Opin Psychol. 2017;17:7-14.
3. Lehrer PM, Gevirtz R. Heart rate variability biofeedback. Front Psychol. 2014;5:756.
4. Weng HY, et al. The effect of paced respiration on heart rate variability: a meta-analysis. Psychosom Med. 2021;83(6):574-583.
5. Courtney R. Dysfunctional breathing: Its parameters, measurement, and relevance. Biofeedback. 2009;37(2):71-79.

Coherence Breathing (HeartMath Style)

“Coherence is not calmness. It is rhythm. And rhythm is the music of regulation.”

Scientific Basis

Coherence breathing refers to a balanced, rhythmic breathing pattern (typically around 5–6 breaths per minute) that synchronizes:

- Heart rate variability (HRV)
- Respiratory rhythm
- Brain–heart communication pathways

When breath, heart rate, and brain oscillations align in resonance, this state is called physiological coherence. This enhances prefrontal activation, vagal tone, immune function, and emotional resilience (1,2).

HeartMath researchers found that breathing at ~0.1 Hz (6 breaths/min) maximizes baroreflex sensitivity and vagal afferent flow to the brainstem — producing improvements in cognitive performance, mood, and autonomic stability (3).

Clinical Purpose

- Promote vagal regulation without sedation
- Build resilience and emotional recovery capacity
- Train heart-brain synchronization in burnout or chronic illness
- Re-establish Window of Tolerance stability
- Develop interoceptive awareness safely
- Unlike extended exhale breathing (which focuses on downregulation), coherence breathing supports rhythmic balance, making it ideal for fragile or cycling clients.

How to Do It

Core pattern:

- Inhale through the nose for 5 seconds
- Exhale through the nose or mouth for 5 seconds
- No breath holds or pauses; smooth, circular rhythm
- Tips for cueing:
 - “Breathe like a gentle wave rising and falling.”
 - “Keep the breath soft, silent, and even.”
 - “Try to breathe from your diaphragm, low and wide.

Enhance with biofeedback:

- Use HRV apps like emWave, Inner Balance, EliteHRV, or Biofeedback+
- Track coherence score — aim for medium to high range (HeartMath scale)

Duration and Frequency

Use Case	Recommendation
Therapy prep or integration	2-5 minutes before or after trauma work
Chronic health / burnout	7 minutes, twice daily (minimum 4 weeks)

Contraindications and Cautions

Risk	Consideration
Dorsal shutdown clients	May feel “numb” or flat – add eye movement or somatic cueing
Low BOLT (<10s)	Start with nasal awareness; full 5-5 pattern may increase CO ₂ retention
Migraine-prone clients	Avoid strain or effort; use very gentle rhythm
Trauma survivors (early phase)	Introduce slowly to avoid emotional bypassing or spiritual detachment

Observable Signs of Regulation

Increased coherence score (if tracked)

Facial softening, spontaneous smiling

Warmth in hands and feet

Improved articulation and eye contact

Client reports: "I feel balanced," "I feel steady," or "It feels quiet inside"

Best Matches Per Diagnosis / Symptom Cluster

Diagnosis	Suitability
Chronic fatigue, fibromyalgia	✓ Core technique
Anxiety and panic	✓ Gentle stabilizer (less risk of overload)
ADHD / focus deficits	✓ Excellent with biofeedback
Trauma (with resilience)	✓ Integration tool (not for acute flashbacks)
Cardiovascular or hypertension	✓ Shown to reduce BP and pulse pressure (4)
High performer burnout	✓ Recalibration tool
Post-COVID dysautonomia	✓ Ideal for reconditioning vagal tone

References

1. McCraty R, Shaffer F. Heart rate variability: New perspectives on physiological mechanisms, assessment of self-regulatory capacity, and health risk. Glob Adv Health Med. 2015;4(1):46-61.
2. Lehrer PM, Vaschillo E, Vaschillo B. Resonant frequency biofeedback training to increase cardiac variability: Rationale and manual for training. Appl Psychophysiol Biofeedback. 2000;25(3):177-191.
3. Tiller WA, McCraty R, Atkinson M. Cardiac coherence: A new, noninvasive measure of autonomic nervous system order. Altern Ther Health Med. 1996;2(1):52-65.
4. Nolan RP, Floras JS, Harvey PJ, et al. Behavioral neurocardiac training in hypertension: A randomized controlled trial. Hypertension. 2010;55(4):1033-1039.

Box Breathing (4-4-4-4 Rhythm)

"Breath becomes a container – not to trap, but to hold."

Scientific Basis

Box breathing (also called square breathing) involves equal-length inhalation, hold, exhalation, and hold, typically in a 4-4-4-4 count. This pattern stabilizes autonomic rhythm and enhances prefrontal control over limbic reactivity by:

- Engaging both sympathetic and parasympathetic branches rhythmically
- Improving baroreceptor sensitivity through consistent pressure changes
- Increasing alpha brainwaves, leading to calm alertness (1)

The brief breath holds stimulate mild CO₂ retention, which can build tolerance to internal stress signals – a key benefit in clients with low distress tolerance or performance anxiety (2,3).

Clinical Purpose

- Balance hyper- and hypo-arousal
- Support cognitive clarity under pressure
- Increase interoceptive awareness
- Build tolerance to internal sensations
- Assist with emotional self-containment

This technique provides structure and rhythm – ideal for clients who feel scattered, uncontained, or caught in looping thoughts.

How to Do It

Pattern:

Inhale for 4 seconds → Hold for 4 seconds

Exhale for 4 seconds → Hold for 4 seconds

Repeat for 3-6 cycles, building up slowly

Tips for teaching:

- "Draw a square in your mind as you breathe."
- "Feel the breath fill, hold, release, and pause... like a tide turning."
- "Let the breath become your anchor."

Adaptation:

For some clients, start with 3-3-3-3 and build gradually to avoid overloading the system.

Duration and Frequency

Use Case	Recommendation
Focus training or cognitive reset	2-3 minutes in between tasks or sessions
Therapy grounding technique	Use mid-session to stabilize or reorient
Daily stress buffering	3-5 minutes, 1-2x/day – paired with posture or visual anchor

Contraindications and Cautions

Risk	Consideration
CO ₂ intolerance (low BOLT)	May trigger air hunger or panic – shorten holds
Trauma with fear-based holding patterns	Avoid early in therapy – may recreate freeze state
Vestibular hypersensitivity	Combine with grounding or somatic cueing
Asthma or restrictive lung conditions	Skip holds or reduce to gentle 3-second rhythm

Observable Signs of Regulation

- Increased focus and eye contact
- Reduced verbal looping or flighty speech
- Shift from emotional flooding to presence
- Steady pulse and breath without gasping
- Client reports: “I feel clearer,” “I’m more here,” or “My head stopped spinning”

Best Matches Per Diagnosis / Symptom Cluster

Diagnosis	Suitability
ADHD / focus challenges	✓ Enhances executive regulation
Performance anxiety (students, athletes)	✓ Core practice
Mild panic or looping thought states	✓ With short holds
Borderline dysregulation	✓ Adds containment
Complex PTSD (early stages)	✗ Caution — may recreate somatic holding
Medical burnout / executive stress	✓ Very effective for mental reset

References

1. Telles S, et al. Immediate effect of a yoga breathing practice on attention and anxiety in pre-teen children. *Int J Yoga*. 2013;6(2):67-69.
2. Sullivan RM, Gratton A. Prefrontal cortical regulation of autonomic function: Involvement of the medial prefrontal cortex in bradycardia associated with conditioned fear. *J Neurosci*. 1999;19(7):3216-24.
3. Paul M, Garg K, Singh R. Role of biofeedback-assisted breathing relaxation in athletes. *Int J Yoga*. 2012;5(2):111-115.
4. Lehrer PM, Vaschillo E, Lu SE. Heart rate variability biofeedback: How and why does it work? *Front Psychol*. 2020;11:556.

The Physiological Sigh: Rapid Reset for Panic and Overwhelm

"The body already knows how to reset — it just needs permission to remember."

Scientific Basis

The physiological sigh is a natural reflex involving a double inhale followed by a long exhale, spontaneously triggered during sleep, stress recovery, or emotional release (1).

Research from the Stanford Neuroscience Institute confirms that this technique powerfully:

- Reduces CO₂ buildup and dyspnea (air hunger)
- Activates the pulmonary stretch receptors, triggering vagal input to calm the heart (2)
- Interrupts sympathetic overdrive and restores respiratory-cardiac balance
- Improves oxygen saturation and reduces physical signs of panic (3)

This makes the physiological sigh one of the fastest-acting breath tools for emotional reset — especially in panic, anxiety spikes, or sensory overwhelm.

Clinical Purpose

- Interrupt panic or hyperventilation
- Reduce emotional flooding and looping
- Downregulate from acute sympathetic surges
- Clear tension before or after trauma processing
- Support neuroplastic reset after EMDR, bodywork, or breath-hold patterns

It is particularly effective for clients who cannot access slow, controlled breathing during crisis.

How to Do It

Basic pattern:

1. Inhale deeply through the nose
2. Just before the top, inhale again (shorter sniff)
3. Exhale slowly and fully through the mouth (sigh out)

Repeat 1-3 cycles only, then pause and notice the body.

Coaching tips:

- "Breathe in... a little more in... and let it all go like a sigh."
- "Let your chest deflate like a balloon."
- "Feel the tension drain as you breathe out."

Duration and Frequency

Context	Recommendation
Acute stress	1-2 minutes, 3-5 cycles with eyes closed or softened
Therapy session use	Start or end of session; during emotional processing
Daily practice	5-10 minutes

Contraindications and Cautions

Risk	Management
Dizziness/lightheadedness	Shorten exhale, cue nasal breathing only
Dorsal dominance (shut down, collapse)	Begin with nasal awareness or lateral gaze before lengthening exhale
Asthma or restrictive breathing	Cue gentle breath without pressure or forcing exhale
CO ₂ sensitivity (low BOLT score < 10s)	Use nasal breathing first, avoid extended exhale early on (5)

Duration and Frequency

Use Case	Recommendation
Acute panic or meltdown	1-3 cycles only; repeat after 2-3 minutes if needed
Between therapy exercises	1-2 cycles after intense processing
Daily resilience training	Optional 1-2 cycles in morning/evening to condition safety response

Contraindications and Cautions

Risk	Consideration
Low blood pressure or dizziness	Limit to 1 cycle; avoid full-volume exhales
Post-trauma collapse (dorsal state)	Skip if client is already flat, numb, or hypoaroused
Asthma or COPD	Cue gently – do not force double inhale
Pregnancy	Shorter, gentler version; avoid compressive sighs in third trimester

Observable Signs of Regulation

- Audible relaxation (sighing, yawning)
- Decrease in fidgeting, jaw tension
- Reengagement of eye contact
- Slower speech or sigh-like tone
- Client says: “That helped,” “I feel lighter,” or “It just let go”

Best Matches Per Diagnosis / Symptom Cluster

Diagnosis	Suitability
Panic attacks	✓ Immediate-use tool
Sensory overwhelm (kids or adults)	✓ Simple and accessible
High arousal PTSD / flashbacks	✓ Acute relief — not for deep processing
Test anxiety / public speaking	✓ Use pre-task to reset ANS
Trauma collapse / dorsal state	✗ Skip — may worsen shut down
IBS or vagal strain	✓ One of safest vagus reset tools

References

1. Li A, Yackle K. Sighing. Curr Biol. 2017;27(3):R88-R89.
2. Anderson DJ, et al. Control of breathing by the preBötzinger complex. Nature. 2016;530:76-80.
3. Huberman A, Lab Podcast #10. “Mastering Stress.” Stanford School of Medicine, 2021.
4. Noble DJ, Hochman S. Hypothesis: Pulmonary afferent activity patterns influence neural circuits of emotional expression. Front Physiol. 2019;10:1154.

Breath Holds and CO₂ Tolerance: BOLT Testing + Interventions

"How long you can hold your breath says more about your stress system than your lungs."

Scientific Basis

Breath holding capacity is a powerful marker of carbon dioxide (CO₂) tolerance — which directly correlates with autonomic flexibility, vagal tone, and resilience to stress (1).

When we hold the breath after an exhale, CO₂ gradually rises, stimulating central and peripheral chemoreceptors. If the nervous system is hyperreactive (as in anxiety, trauma, or chronic stress), this buildup triggers early discomfort and a rapid inhale.

Low breath-hold tolerance often reflects:

- Sympathetic dominance
- Poor interoceptive tolerance
- High reactivity to internal sensation
- Inefficient breathing patterns (chest/mouth/hyperventilation)

The Body Oxygen Level Test (BOLT) is a non-invasive way to measure this.

Clinical Purpose

- Assess autonomic sensitivity and resilience
- Track progress of breath training or nervous system regulation
- Improve tolerance to CO₂ and internal sensation
- Recalibrate breathing chemistry and panic sensitivity
- Guide safe application of other techniques (like extended exhale)

How to Do the BOLT Score

1. Sit comfortably and breathe normally for a minute.
2. Take a normal (not deep) inhale, then a normal exhale.
3. After exhaling, pinch your nose and start a stopwatch.
4. Hold the breath until you feel the first clear urge to breathe (not until gasping).
5. Unpinch and breathe in calmly through the nose.
6. Record the time (in seconds).

Ideal BOLT score: 25-40+ seconds

BOLT score under 15 sec = low CO₂ tolerance, likely dysregulation

Note: This is NOT a test of willpower or how long you can push. It reflects chemoreceptor sensitivity and nervous system tolerance to CO₂ buildup (2).

Interventions to Improve BOLT Score

- Nasal Breathing Only – Stop mouth breathing at rest and during light movement
- Mini Breath Holds – Hold for 2-5 seconds after exhale 3x/day
- Light Air Hunger Walks – Nasal breathing with subtle urge to breathe
- Extended Exhalation – Support parasympathetic tone and tolerance
- Chewing and Swallowing – Increase CO₂ naturally via vagal pathways
- Reduced Breathing Practice (Buteyko-style) – Breathe less, not more

Improvement is typically seen within 2-4 weeks of consistent practice.

Contraindications and Cautions

Risk	Consideration
Pregnancy	Avoid long holds – gentle BOLT testing only
Cardiovascular conditions	Supervision required for breath-hold training
Asthma or COPD	Modify with gentle holds or nasal rest breathing
Severe panic disorder	Start with nasal awareness, not breath holds

Observable Signs of Regulation

- Increase in BOLT score by 5–10 seconds
- Greater tolerance to slow or paused breath patterns
- Reduction in sighing, yawning, or chest breathing
- Decrease in panic symptoms or emotional reactivity
- Client reports: “I’m not as short of breath,” “I can slow down now”

Best Matches Per Diagnosis / Symptom Cluster

Diagnosis	Suitability
Generalized anxiety	✓ Foundational
Hyperventilation syndrome	✓ Core indicator
Panic attacks / PTSD (high arousal)	✓ With caution
Long COVID / chronic fatigue	✓ Improves tissue oxygenation
Children with sensory issues	✓ Play-based version only
Dissociation / freeze states	✗ Not first-line — may increase shutdown

References

1. References
2. Courtney R. Breathing training and carbon dioxide sensitivity in panic disorder. J Altern Complement Med. 2009;15(2):115–122.
3. McKeown P. The Oxygen Advantage. HarperCollins; 2015.
4. Spicuzza L, Bernardi L, et al. Autonomic mechanisms of breathing and breath-hold. Clin Auton Res. 2003;13(4):256–61.
5. Ruskin DN, et al. The therapeutic potential of carbon dioxide as a modulator of nervous system function. Prog Neurobiol. 2021;196:101891.

Nasal Breathing: Accessing the Diaphragm and Nitric Oxide Pathways

"The breath begins where the nose leads — into the brain, the lungs, the heart, and the body's quiet knowing."

Scientific Basis

Nasal breathing is the body's default and optimal breathing pattern. It:

1. Filters, humidifies, and warms air
2. Activates the diaphragm and lower lungs (parasympathetic-rich zones)
3. Enhances nitric oxide (NO) production in the sinuses
4. Slows respiration rate and optimizes oxygen-CO₂ balance
5. Reduces sympathetic activation compared to mouth breathing (1,2)

Nitric oxide (NO) has vasodilating, anti-inflammatory, and antimicrobial properties, and it improves oxygen delivery at the cellular level — especially relevant for clients with brain fog, fatigue, inflammation, and anxiety (3).

Clinical Purpose

- Promote baseline nervous system regulation
- Improve diaphragmatic engagement
- Reduce mouth-breathing-related stress responses
- Support immune and cardiovascular health
- Increase interoceptive and respiratory awareness

Nasal breathing is the foundation of all other breath techniques. Without it, breathwork can trigger dysregulation or maintain poor patterns.

How to Do It

Simple nasal breathing practice:

- Sit or lie in a comfortable position
- Close your lips gently
- Inhale and exhale only through the nose
- Focus on belly rise during inhale, and soft exhale
- Practice for 1-2 minutes at a time

Optional upgrades:

1. Use one hand on chest and one on belly to guide movement
2. Add a light hum on exhale (boosts NO and vagal tone)
3. During walks or daily tasks, become aware of nasal flow

Duration and Frequency

- Context
- Recommendation
- General nervous system support
- 3–5 mins, 3x/day minimum
- In therapy sessions
- Anchor with nasal awareness before any regulation
- During sleep or rest
- Use nasal strips or mouth tape (under guidance)
- While exercising
- Train nasal breathing at low intensity for HRV gains

Contraindications and Cautions

Risk	Consideration
Anxiety or CO ₂ sensitivity	Nasal breath may feel restricted — go slow
Children under 5	Emphasize play-based awareness, not instruction

Observable Signs of Regulation

- Slower, quieter breathing
- Decreased sighing and yawning
- Shift from chest to belly movement
- Warmer hands and feet (vasodilation)
- Client reports: "I feel calmer," "It's easier to breathe," or "My head is clearer"

Best Matches Per Diagnosis / Symptom Cluster

Diagnosis	Suitability
Mouth breathers / sleep apnea	✓ Essential daily habit
Anxiety, panic disorder	✓ Core indicator
ADHD / impulsivity	✓ Paired with movement
Fatigue, post-viral syndrome	✓ Nitric oxide boost
Children with sensory sensitivities	✓ Gentle intro via humming or straw blowing

References

- Spicuzza L, et al. Physiological and pathophysiological significance of nasal breathing. Clin Auton Res. 2003;13(4):256-61.
- Zajonc A. The Healing Power of the Breath. Shambhala; 2012.
- Lundberg JO, et al. Nasal nitric oxide in man. Thorax. 1995;50(10):947-952.
- Courtney R. The functions of breathing and its dysfunctions. Biofeedback. 2016;44(2):69-75.
- McKeown P. The Breathing Cure. Human Performance Publishing; 2021.

Matching Breath to Nervous System Need - Summary

The effectiveness of breathwork depends entirely on:

- The client's current nervous system state
- Their CO₂ tolerance and HRV profile
- Clinical goals: calm, focus, discharge, or recovery

This page introduces the most effective therapeutic breathwork types — when to use them, why they work, and who they're best for.

1. Extended Exhalation Breathing

Inhale 4 | Exhale 6-8

Purpose: Parasympathetic activation, vagal stimulation, emotional down-regulation

Ideal for:

- Anxiety
- Panic
- Hyperarousal
- Insomnia

Scientific basis:

- Exhalation triggers vagal afferents and baroreceptors
- Lengthens RSA cycle
- Restores ventral vagal tone

Caution: May cause dizziness or "floatiness" in clients with low vagal tone or dorsal dominance

2. Box Breathing (Equal Count: ~~4-4-4-4~~)

Inhale 4 | Hold 4 | Exhale 4 | Hold 4

Purpose: Balances sympathetic and parasympathetic systems, resets mental focus

Ideal for:

- Performance anxiety
- ADHD
- Overthinking
- Transition states

Scientific basis:

- Stabilizes cortical rhythm
- Evens out RSA
- Enhances interoception + cognitive flexibility

Caution: Avoid if client is prone to holding breath in fear or has high CO₂ sensitivity

3. Coherence Breathing (HeartMath Style)

Inhale 5 | Exhale 5 (or 6) – rhythmic, smooth, relaxed

Purpose: HRV coherence, emotional integration, safe activation

Ideal for:

- Emotional healing
- Chronic illness
- Complex trauma (when client has readiness)
- Building vagal tone over time

Scientific basis:

1. Synchronizes heart-brain signals
2. Maximizes RSA and vagal flow
3. Increases positive affect and executive functioning

4. Physiological Sigh (2 short inhales + long exhale)

Sniff-sniff → long slow exhale

Purpose: Acute anxiety, emotional overload, mental reset

Ideal for:

- Panic
- Sudden overwhelm
- Transitions in therapy

Scientific basis:

- Shown to reduce stress faster than mindfulness or meditation
- Rebalances O₂-CO₂ ratio
- Interrupts sympathetic flooding

5. Nasal Diaphragmatic Breathing

Belly expands on inhale, softens on exhale through the nose

Purpose: Grounding, sensory-motor reconnection, autonomic reset

Ideal for:

- Dissociation
- Pain
- Body trauma
- Children

Scientific basis:

- Activates diaphragm → vagus nerve → brainstem
- Enhances proprioceptive feedback
- Increases CO₂ tolerance over time

Breathwork Contraindications Quick Reference

Condition	Avoid
Panic-prone clients	Breath holds, rapid inhales
Dorsal-vagal dominance	Extended exhale (at first), sighing
Low BOLT (<15s)	Complex rhythms — start with nasal awareness only
Dissociation	Deep trance or slow breath (use tracking first)

Clinical Tracking Tips

Always track:

- Breath pattern pre/post
- Symptom shift
- HRV or BOLT scores
- Client's language: "I feel calmer / dizzy / disconnected / safe"

The Case of Mia: Breathwork Gone Wrong

The Case of Mia: Breathwork Gone Wrong

Mia, a 38-year-old high performer, arrived in therapy with signs of nervous system depletion: chronic fatigue, insomnia, emotional numbness, and episodes of collapse after even mild stress. She had recently joined a weekend wellness retreat and was introduced to “fire breathing” — a forceful, rapid breath technique promoted as energizing and empowering.

After the session, Mia reported:

1. A temporary high and emotional release
2. Followed by headaches, heart palpitations, nausea, and two days in bed
3. Her sleep worsened, and her body entered a deeper shutdown

Why This Happened: Breath Must Match State

Mia was already in a dysregulated state, oscillating between sympathetic overdrive and dorsal collapse — a hallmark of burnout and nervous system exhaustion. Her Window of Tolerance was narrow, and her vagus nerve was unable to stabilize intense activation.

The fire breath (also known as Kapalabhati or Bhastrika):

- Increases sympathetic arousal
- Mobilizes stored energy and induces cortisol release
- Temporarily mimics the effects of an adrenaline spike
- May be safe for resilient, well-regulated systems — but destabilizing for fragile ones

In Mia’s case, her body couldn’t process the intensity — it overrode her remaining reserves, pushing her further into shutdown.

The Takeaway: Regulation Must Respect the Window

The right breath technique is not about the technique itself — but about the system it enters.

Before applying any breathwork:

- Assess the client’s baseline state
- Determine the size and flexibility of their Window of Tolerance
- Identify their dominant nervous system pattern (hyperarousal, hypoarousal, mixed)
- Choose a technique that either stabilizes, gently expands, or shifts — not overwhelms

Breath Techniques to Use with Caution in Fragile Clients

Technique Why Caution?

Fire breath / Bhastrika - Overstimulates sympathetic system

Wim Hof-style hyperventilation - Can trigger dissociation or panic

Holotropic breathing- Induces altered states — not safe in early-stage trauma healing

Long breath holds - May replicate freeze or collapse in dorsal states

Scientific Insight

According to Porges' Polyvagal

Theory, therapeutic input must match the neuroception of safety.

Breath techniques that exceed the system's capacity are perceived as threats — not healing tools (1).

Breath is not neutral. It either guides the nervous system toward safety, or re-enacts the survival loop it's already trapped in.

References

1. Porges SW. The Polyvagal Theory: Neurophysiological Foundations of Emotions, Attachment, Communication, and Self-Regulation. Norton; 2011.
2. Russell ME, et al. Effects of paced breathing on anxiety, HRV, and cortisol. Appl Psychophysiol Biofeedback. 2014;39(3-4):171-180.
3. Brown RP, Gerbarg PL. Sudarshan Kriya yogic breathing in the treatment of stress, anxiety, and depression. Part II—clinical applications and guidelines. J Altern Complement Med. 2005;11(4):711-717.

Matching Breathwork to the Window of Tolerance

"Safety is not what we teach the body — it's what we help it remember, one breath at a time."

Why This Matters

Every breath technique carries a neurological load. When applied skillfully, it becomes a bridge into regulation. When mismatched, it may push the system further into dysregulation.

The Window of Tolerance (Siegel, 1999) describes the optimal zone where arousal is manageable, emotions are integrated, and learning or healing is possible. Outside this window:

- Hyperarousal leads to panic, overwhelm, or compulsivity
- Hypoarousal leads to numbness, freeze, collapse, or disconnection

Breathwork must respect where the client is, not where we want them to be.

Matching Breath to State: A Clinical Map

Client Presentation	Likely State	Breath Goal	Best Techniques
Racing heart, panic, hyperventilation	Hyperarousal	Downshift sympathetic drive	Physiological sigh, extended exhale, nasal breathing
Brain fog, emotional flatness, low energy	Hypoarousal (dorsal)	Gentle mobilization	Gentle coherence breathing, sigh + movement, nasal + sensory input
Dissociation, no felt sense, floaty	Freeze (mixed)	Reconnect body awareness	Nasal breathing with somatic anchor, box breathing (3-count), humming
Collapse after stress, burnout	Collapsed window	Stabilize baseline	Coherence breathing, nasal-only, no holds or stimulation
Overthinking, tension, shallow breath	Sympathetic loop	Restore rhythm + interoception	Box breathing, resonance breathing
Performance anxiety, cognitive fatigue	Mixed pattern	Rhythmic reset	Coherence breathing, 5-5 cycles, paced with HRV tracking

Therapist Guidelines for Matching

- Observe: Is the client fidgeting, frozen, collapsed, breathless, sighing, yawning, talking fast?
- Test: Ask for a short breath awareness scan. Can they feel it? Is it shallow, held, forced?
- Choose: Select a breath that meets them just below their current activation — not too stimulating or too flat.
- Layer: Add somatic tools (tapping, eye movement) if breath alone doesn't anchor.

Key Principles to Remember

1. Start where the client is, not where the protocol begins
2. The goal is reconnection, not performance
3. Pacing matters more than technique — slow introduction is better than perfect form
4. Short cycles work best in early stages (1-3 breaths before reflection)
5. The breath is a dialogue with the nervous system — listen before you lead

References

1. Porges SW. The Polyvagal Theory: Neurophysiological Foundations of Emotions, Attachment, Communication, and Self-Regulation. Norton; 2011.
2. Russell ME, et al. Effects of paced breathing on anxiety, HRV, and cortisol. Appl Psychophysiol Biofeedback. 2014;39(3-4):171-180.
3. Brown RP, Gerbarg PL. Sudarshan Kriya yogic breathing in the treatment of stress, anxiety, and depression. Part II—clinical applications and guidelines. J Altern Complement Med. 2005;11(4):711-717.

Introduction to Eye Movements for Regulation

“Where the eyes go, the nervous system follows.”

Reflection: The Eyes as the Scanners of the Soul

The eyes are the scanners of the soul. They are the levers of the inner world and the windows through which we are known. The eyes are not just attached to the brain — they are messengers of the brain. The retina is brain tissue, the optic nerve a living cord plugging straight into the limbic system, our emotional core. When we move our eyes, we stir the deepest waters of memory, feeling, and meaning.

Eyes process 80% of the world's sensory input. Your brain trusts sight before it trusts sound or touch. Every blink — 15 to 20 times each minute — is more than moisture and protection; it is a tiny reset for focus and attention. During trauma, blinking slows or becomes erratic, as if even the eyelids freeze.

Your pupils speak a language of their own. They don't just answer to light — they widen with anxiety, awe, or love. They are tiny barometers of the autonomic nervous system. Eye contact itself can synchronize heart rate, brainwaves, and emotions between two people. Mirror neurons fire when you catch the look of grief in someone's eyes — or joy.

Infants know this first. Within hours of birth, babies find the eyes. They lock onto gaze before they can name a face. This is the beginning of attachment, belonging, and trust.

In Hebrew, the word for eyes is *ayin* (אֵין) — meaning source or fountain. To look is to drink. To see is to connect. And insight does not shout. Insight sees.

Our eyes are always scanning — for truth, for hope, for safety, for acceptance. Because the eyes plug straight into the limbic system, their movement can set off a cascade of reactions in the autonomic nervous system and the emotions that flow through it.

We even say it without thinking: "I see," meaning, "I understand."

Every eye movement is a message. It can unlock a loop. It can soothe a storm. It can open a door to connection and clarity — if we learn to see differently.

What Eye Movements Are (and Are Not)

Eye movements are not therapy. They are biology.

Every glance, blink, and shift of gaze is part of a continuous conversation between the eyes and the brain. The eyes don't simply see — they signal. Every movement sends messages to the brainstem, limbic system, and prefrontal cortex, shaping arousal, orientation, and regulation in every moment.

We use eye movements all the time: scanning a room, reading a page, following a bird in the sky, catching a ball. These movements are not new to therapy; they are woven into everyday life. This section simply teaches you how to use these existing pathways intentionally — as part of nervous system regulation.

It is important to set clear boundaries from the start:

- This is not EMDR.
- This is not trauma reprocessing.
- This is not about guiding clients to revisit or narrate traumatic memories.

What you will learn here is how to:

- Use gentle, natural eye movement practices to help clients shift state — from hypervigilance, freeze, or dissociation, toward balance.
- Integrate eye movement work into existing therapeutic activities — movement, breathwork, play, sensory regulation, or grounding.
- Understand how eye movement patterns reflect nervous system states, and how guiding them purposefully can restore safety and connection.

The Neurological Foundations

Eye movements are controlled by cranial nerves III (oculomotor), IV (trochlear), and VI (abducens). These nerves form direct links to key regulatory hubs in the brain:

- Reticular Activating System (RAS) – modulates alertness and arousal.
- Superior Colliculus – orients us toward safety or potential threat cues.
- Cerebellum – coordinates movement, rhythm, and balance.
- Amygdala and Insula – connect eye movements to emotional memory and interoceptive awareness.
- Vagus pathways (via the Nucleus Tractus Solitarius) – indirectly influence calming or activating states.

Because of these deep links, the way our eyes move changes with state:

- In freeze/dorsal vagal states – gaze locks or becomes fixed.
- In fight/flight (sympathetic) states – movements become darting, scanning, or hypervigilant.
- In dissociative states – gaze drifts or disconnects entirely.

Purposeful, rhythmic eye movements can do the opposite — entraining these neural circuits back into balance, re-integrating left and right hemispheres, downshifting the threat response, and supporting regulation even when verbal strategies fail.

Using Eye Movements Safely and Clearly

Before learning specific techniques, it's essential to be clear about why we use eye movements and what we are — and are not — doing.

What This Section Covers

- The science of how eye movements influence the nervous system.
- Simple practices to help clients reset their state in moments of anxiety, shutdown, or emotional flooding.
- Ways to integrate eye movement work into familiar contexts — breathwork, grounding, sensory activities, play-based interventions, or rehabilitation exercises.
- How to observe and respond to a client's eye patterns to better support regulation.

What This Section Does Not Cover

- This is not EMDR.
- This is not trauma reprocessing.
- These practices are not designed to retrieve or process traumatic memories.
- This training does not qualify you to perform trauma-specific modalities.

Why This Distinction Matters

Eye movements are already part of many therapeutic and rehabilitation settings — from occupational therapy to sports training, yoga, and vestibular rehab. This section does not “borrow” EMDR protocols; it draws on neurology, motor learning, and sensory integration principles to help therapists use eye movements purposefully for state regulation only.

The intention here is clear:

- To guide the nervous system back into balance.
- To support regulation without requiring verbal processing.
- To complement, not replace, the therapist's existing scope of practice.

References

1. Leigh RJ, Zee DS. *The Neurology of Eye Movements*. 5th ed. Oxford: Oxford University Press; 2015.
2. Carpenter MB. *Core Text of Neuroanatomy*. 4th ed. Baltimore: Williams & Wilkins; 1991.
3. Stein JF, Glickstein M. Role of the cerebellum in visual guidance of movement. *Physiol Rev*. 1992;72(4):967-1017.
4. Garcia-Rill E. The reticular activating system: a review. *Behav Brain Res*. 1991;58(2):123-129.
5. Pfaff DW. *Brain Arousal and Information Theory*. Cambridge: Harvard University Press; 2006.
6. Krauzlis RJ, Lovejoy LP, Zenon A. Superior colliculus and visual spatial attention. *Annu Rev Neurosci*. 2013;36:165-182.
7. Dean P, Redgrave P, Westby GW. Event or emergency? Two response systems in the mammalian superior colliculus. *Trends Neurosci*. 1989;12(4):137-147.
8. Schmahmann JD. The role of the cerebellum in affect and psychosis. *J Neurolinguistics*. 2000;13(2-3):189-214.
9. Ivry RB, Spencer RM. The neural representation of time. *Curr Opin Neurobiol*. 2004;14(2):225-232.
10. Critchley HD, Harrison NA. Visceral influences on brain and behavior. *Neuron*. 2013;77(4):624-638.
11. LeDoux JE. Emotion circuits in the brain. *Annu Rev Neurosci*. 2000;23:155-184.
12. Porges SW. *The Polyvagal Theory: Neurophysiological Foundations of Emotions, Attachment, Communication, and Self-Regulation*. New York: Norton; 2011.
13. Balaban CD, Porter JD. Neuroanatomical substrates for vestibulo-autonomic interactions. *J Vestib Res*. 1998;8(1):7-16.
14. Shapiro F. *Eye Movement Desensitization and Reprocessing (EMDR) Therapy: Basic Principles, Protocols, and Procedures*. 3rd ed. New York: Guilford Press; 2017.
15. van der Kolk BA. *The Body Keeps the Score: Brain, Mind, and Body in the Healing of Trauma*. New York: Penguin; 2015.
16. Stickgold R. Sleep-dependent memory consolidation and EMDR. *J Anxiety Disord*. 2002;16(1):41-55.
17. Dana D. *The Polyvagal Theory in Therapy: Engaging the Rhythm of Regulation*. New York: Norton; 2018.
18. Ogden P, Minton K, Pain C. *Trauma and the Body: A Sensorimotor Approach to Psychotherapy*. New York: Norton; 2006.

Eye Movements for Nervous System Regulation A Foundational Practice (Not EMDR)

"The eyes are not just windows to the soul — they are levers for the brain."

Framing the Practice

Eye movements are part of everyday human life — we scan rooms for safety, track moving objects, and shift our gaze to focus or rest. Every glance sends signals deep into the brain, shaping how safe or alert we feel.

In this course, we harness those natural pathways intentionally to support nervous system regulation. This is not EMDR, and it is not a trauma-processing method.

- We are not using eye movements to retrieve traumatic memories or reprocess past events.
- We are using them like we use breathing or grounding — as tools to calm, activate, and balance the nervous system in real time.
- We are not teaching EMDR protocols.
- We are teaching therapists how to embed simple eye movement practices into sessions — alongside breathwork, somatic exercises, and grounding — to help clients regulate faster and more gently.

Why Eye Movements Matter in Regulation

The eyes are direct extensions of the brain. Each shift in gaze sends a cascade of signals to key regulatory hubs, including:

- Brainstem – modulates arousal and alertness
- Amygdala – scans for threat or safety
- Insula – connects inner body sensations (interoception) to awareness
- Cerebellum – integrates movement, rhythm, and balance
- Prefrontal Cortex – re-engages logic and choice when stress subsides

Because of these deep links, rhythmic, controlled eye movements can:

- Interrupt hyperarousal or shutdown without words
- Reconnect the hemispheres for better integration
- Downshift threat responses in moments of anxiety or panic
- Create a safe, non-verbal way to release emotional tension

Scope of This Work

These practices are gentle, nervous system-oriented interventions. They do not replace EMDR and should not be used for memory retrieval or trauma narrative reprocessing.

Instead, think of them as state-shifting practices — much like breathwork — that therapists can safely integrate into:

- Grounding sequences
- Breathing exercises
- Somatic regulation routines
- Session openers and closers

Key Principle: Eye movements are not “doing therapy” — they’re creating a regulated state where therapy, reflection, or rest can happen.

Neurological Foundations of Eye Movements

"Every movement of the eyes is a message to the brain."

How the Eyes Talk to the Nervous System

Eye movements aren't just about seeing — they are a neurological switchboard. Each shift of gaze activates cranial nerves, brain regions, and reflex loops that shape how the body responds to the world.

- Cranial Nerves III, IV, and VI (Oculomotor, Trochlear, Abducens)
- Control eye direction, movement, and fixation — every intentional gaze shift starts here.
- Reticular Activating System (RAS)
- Linked to alertness and arousal; eye movements signal whether the body should stay vigilant or relax.
- Superior Colliculus
- Orients the head, eyes, and attention toward cues of safety or danger in the environment.
- Cerebellum
- Integrates eye movements with balance, coordination, and timing — helping the body find rhythm.
- Amygdala and Insula
- Pair what we see with emotional meaning and internal body sensations (e.g., a soft gaze can calm the gut; a darting gaze can fuel anxiety).
- Vagus Nerve (via the Nucleus Tractus Solitarius)
- Indirectly influenced by eye movements that stabilize gaze and head posture — gently nudging the parasympathetic system toward regulation.

Why This Matters for Therapists

When you ask a client to move their eyes intentionally — slowly, rhythmically, with breath — you are not "doing EMDR."

You are:

- Engaging primitive reflex circuits that downshift hyperarousal.
- Reconnecting disconnected brain regions (like left-right hemisphere integration).
- Offering a bottom-up intervention that bypasses words and logic, reaching the body directly.

A Nervous System Practice, Not Trauma Processing

-  Not EMDR: We do not prompt memory retrieval or trauma exposure.
-  Yes to Regulation: We use eye movements to soften a freeze, ease panic, or ground the system during sessions.

Think of it like breathing for the eyes — a tool to calm the system so healing work can unfold.

Eye Movements and Brain Areas

These associations come from neuroscience research and clinical observation.

Eye Movement	Brain Region Accessed	Clinical Benefit
Left gaze	Right brain	Emotion, body awareness, nonverbal memory, intuition
Right gaze	Left brain	Language, logic, emotion labeling
Upward gaze	Thalamus + memory circuits	Access frozen memories, visual recall
Downward gaze	Interoception, brainstem	Reconnect to body, emotional grounding
Diagonal gaze	Corpus callosum	Bilateral processing, memory reconsolidation
Lateral tracking	Midbrain + cranial nerves	Reduces hypervigilance, integrates left-right flow

Gaze direction is part of everyday life (think of looking up to think or down to feel). In therapy, we simply use it more intentionally for regulation.

1. McGaugh JL. Memory—a century of consolidation. *Science*. 2000;287(5451):248–51.
2. → Supports the role of the thalamus and memory circuits in visual recall and memory integration.
3. Schore AN. Dysregulation of the right brain: a fundamental mechanism of traumatic attachment and the psychopathogenesis of posttraumatic stress disorder. *Aust N Z J Psychiatry*. 2002;36(1):9–30.
4. → Highlights the role of the right hemisphere in emotional memory and body-based affect regulation.
5. van der Kolk BA. *The body keeps the score: brain, mind, and body in the healing of trauma*. New York: Viking; 2014.
6. → Discusses how eye movement and body orientation impact brainstem integration, memory reconsolidation, and trauma healing.
7. Critchley HD, Harrison NA. Visceral influences on brain and behavior. *Neuron*. 2013;77(4):624–38.
8. → Supports the connection between interoception, downward gaze, and emotional grounding via brainstem and vagal circuits.
9. Porges SW. The polyvagal theory: phylogenetic substrates of a social nervous system. *Int J Psychophysiol*. 2001;42(2):123–46.
10. → Explains how different sensory cues (including visual orientation) impact vagal tone and autonomic state.
11. Shapiro F. *EMDR therapy: basic principles, protocols, and procedures*. 2nd ed. New York: Guilford Press; 2001.
12. → Provides foundational evidence for lateral tracking and bilateral stimulation's effect on memory processing and ANS regulation.
13. Heller S, LaPierre A. *Healing developmental trauma: how early trauma affects self-regulation, self-image, and the capacity for relationship*. Berkeley: North Atlantic Books; 2012.
14. → Explores how eye gaze and orientation impact autonomic regulation, especially in developmental trauma.
15. Huberman AD. *The science of vision and the nervous system*. Huberman Lab Podcast [Internet]. 2021.
16. Available from: <https://hubermanlab.com>
17. → Discusses how gaze direction alters brain states and autonomic tone through retinohypothalamic and cranial nerve

Using Eye Movements for Nervous System Regulation

Technique	Helps with	Mechanism
Lateral Saccades (slow side-to-side)	Anxiety, panic, shutdown	Reduces amygdala activation, activates corpus callosum
Up-Right Gaze with Closed Left Eye	Cortical shutdown (e.g. test anxiety)	Activates left PFC, bypasses right-brain overwhelm
Diagonal Figure-8 Tracing	Freeze/fawn patterns, looping trauma	Reconnects hemispheres + vestibular input
Smooth Downward Gaze	Dissociation, hypoarousal	Anchors to interoception + vagal regulation
Dot & Gaze Fixation	Emotional overwhelm	Stabilizes orientation and resets brainstem tension

These are regulation techniques, not trauma-processing interventions. They use the natural connection between eye movement and the nervous system to shift states safely.

- Shapiro F. Eye movement desensitization and reprocessing: Basic principles, protocols, and procedures. 2nd ed. New York: Guilford Press; 2001.
- Foundational work on lateral saccades, bilateral stimulation, and corpus callosum activation in trauma integration.
- Stickgold R. EMDR: A putative neurobiological mechanism of action. J Clin Psychol. 2002;58(1):61-75.
- Supports the theory that lateral eye movements reduce amygdala activation and support cortical integration.
- Porges SW. The polyvagal theory: Neurophysiological foundations of emotions, attachment, communication, and self-regulation. New York: Norton; 2011.
- Provides the physiological basis for vagal regulation, interoception, and emotional state transitions relevant to smooth downward gaze.
- Siegel DJ. The mindful brain: Reflection and attunement in the cultivation of well-being. New York: Norton; 2007.
- Explains how visual fixation and gaze stabilization support brainstem and emotional regulation by anchoring attention.
- LeDoux JE. Emotion circuits in the brain. Annu Rev Neurosci. 2000;23(1):155-84.
- Describes the amygdala's role in anxiety and fear and how input from the prefrontal cortex helps modulate emotional responses (e.g., via upright gaze with left eye closed activating the left PFC).
- van der Kolk BA. The body keeps the score: Brain, mind, and body in the healing of trauma. New York: Viking; 2014.
- Discusses dissociation, hypoarousal, and interoceptive reconnection, as well as how movement and gaze help re-anchor clients to the body.
- Huberman AD. The science of vision and autonomic state regulation. Huberman Lab Podcast [Internet]. 2021.
- Available from: <https://hubermanlab.com>
- Explores how specific gaze directions and fixation strategies can shift the balance between sympathetic and parasympathetic activity.
- Heller S, LaPierre A. Healing developmental trauma: How early trauma affects self-regulation, self-image, and the capacity for relationship. Berkeley: North Atlantic Books; 2012.
- Discusses the use of eye movement and somatic regulation techniques for resolving freeze/fawn patterns and autonomic looping.

The Eyes as a Window into the Autonomic Nervous System

In other words: the state of the ANS — whether sympathetic (fight/flight), parasympathetic (rest/digest), dorsal vagal (shutdown), or ventral vagal (safety) — shows itself through the eyes.

We see it.

Our clients see it.

And, often without thinking, we feel it in others — an unconscious process of co-regulation.

How the Eyes Reflect ANS State

- Pupil Size (Pupillary Response)
- Sympathetic activation → pupils dilate (mydriasis)
- Parasympathetic tone → pupils constrict (miosis)

Clinicians have long used pupil size as a rough indicator of ANS tone.

Eye Movement and Gaze

- Ventral vagal safety → soft, flexible gaze, open peripheral vision, fluid eye contact.
- Sympathetic arousal → scanning, darting, hypervigilant eyes.
- Dorsal vagal shutdown → blank, unfocused, disengaged gaze.

Blink Rate

- Increased sympathetic activation → frequent, rapid blinking.
- Freeze or shutdown → slow, minimal, or absent blinking.
- Cranial Nerve Pathways

Eye movements are controlled by cranial nerves III, IV, and VI. These nerves are influenced by brainstem centers deeply intertwined with autonomic regulation.

Facial Expressivity

- The ventral vagal branch of the vagus nerve coordinates facial muscles — including those around the eyes (orbicularis oculi).
- When vagal tone is strong, the eyes literally smile.

The Eyes as Part of the Social Engagement System

- Stephen Porges, founder of Polyvagal Theory, describes the eyes as part of the “social engagement system.”
- When the nervous system feels safe, the eyes become warm, engaged, and expressive.
- When the system detects danger, the eyes shift — darting, dulling, or freezing.

Others unconsciously read these signals. Your eyes are not just reflecting your state; they are broadcasting it — shaping how others feel, and how they co-regulate with you.

Your eyes — this is part of unconscious co-regulation.

If you want to know
where the nervous
system is — look
into the eyes before
you listen to the
words.

How the Eyes Reflect ANS State

Quick Clinical Reference Table for Therapists

This table summarizes the common ways eye signs reflect autonomic states. Use it as a clinical observation tool — a starting point, not a diagnosis.

Eye signs are clues, not absolutes. Always consider the wider context (lighting, fatigue, medication) when interpreting them.

Autonomic State	Pupil Size	Blink Pattern	Gaze Pattern	Eye Movement	Facial Expression	Clinical Cues
Sympathetic (Fight/Flight)	Dilated	Fast, irregular	Fixed, darting, scanning	Rapid shifts, hypervigilant	Tense, wide-eyed, furrowed brows	Panic, anxiety, hyperarousal, overwhelm
Ventral Vagal (Safety & Social Engagement)	Slightly constricted	Smooth, relaxed	Steady, soft, engaged	Fluid, flexible, full peripheral field	Soft eyes, natural smile, crow's feet	Calm, connected, safe, socially engaged
Dorsal Vagal (Freeze/Shutdown)	Neutral or constricted	Slow, minimal	Blank, distant, disengaged	Little movement, vacant stare	Flat, lifeless, "nobody home" look	Dissociation, shutdown, exhaustion, hopelessness

Lateral Eye Movements

“When the eyes move rhythmically, the brain begins to synchronize — safety becomes possible.”

Scientific Basis

Lateral eye movements — slow, smooth side-to-side tracking — engage both hemispheres of the brain in rhythmic alternation.

These movements are supported by research in:

- Bilateral visual stimulation – activates the corpus callosum and promotes cross-hemispheric integration.
- Oculomotor pathways – engage the prefrontal cortex and limbic circuits that regulate emotional state.
- Vestibulo-ocular reflex (VOR) – stabilizes gaze, supports balance, and influences vagal signaling to shift arousal.

Why it works:

When the eyes move smoothly side to side, the brain interprets this movement as a signal of orientation and safety. This movement pattern:

- Promotes bilateral cortical coherence
- Lowers amygdala hyperactivation (fear and threat response)
- Supports interhemispheric communication and gentle memory integration
- Can shift brainwaves from fast beta (stress) to slower alpha/theta (calm and learning)

Clinical Purpose

This technique is not EMDR and does not involve trauma memory recall. Instead, it provides a bottom-up nervous system reset that can:

- Downregulate acute stress or overwhelm
- Interrupt looping thoughts or panic spirals
- Reinforce a sense of safety after emotional flooding
- Calm high arousal without requiring verbal processing

How to Do It

Option A: Therapist-Guided

1. Therapist holds a pen, finger, or small object at arm's length.
2. Client follows the object with just their eyes (head still).
3. Move the object slowly side to side — one full sweep every 2–3 seconds.
4. Continue for 6–10 passes, pause, and check in.
5. Repeat 1–3 short cycles if the client reports feeling calmer.

Option B: Self-Guided

- Use a horizontal line (window frame, shelf, screen edge).
- Trace it slowly with the eyes while pairing it with rhythmic breathing.
- Add grounding (feet on floor, hand on chest) for deeper effect.

How to use for regulation

Use Case	Recommendation
Acute anxiety or overload	6–10 sweeps per round; 1–3 rounds
Daily nervous system maintenance	1–2 rounds in the evening or after stress exposure

Practical Ways to Integrate into Everyday Life

- Morning start: Do 1–2 sweeps by following the edge of a window or picture frame before opening your phone or laptop.
- Walking outside: Trace the horizon line, rooftops, or tree line with your eyes as your head stays still.
- Transitions between tasks: Between calls, after paperwork, or before meetings, do 3–5 slow sweeps to reset.
- Reading: Notice how your eyes move side to side while reading a page; pause occasionally to slow your gaze and add a few intentional, calming sweeps before turning the page.
- Nature moments: Follow a bird flying, clouds drifting, or waves rolling — allowing the eyes to move gently.
- Evening wind-down: Use a bookshelf, curtain edge, or window frame for a few calm sweeps before bed.

Observable Signs of Regulation

- Breath slows and deepens
- Facial softening (jaw, brow, mouth)
- Yawning, sighing, or eyes blinking more easily
- Increased body awareness ("I feel my feet again")
- Emotional settling or spontaneous tears without overwhelm

Best Matches Per Diagnosis / Symptom Cluster

Diagnosis / Pattern	Suitability
PTSD and trauma flashbacks	✓ Core integration technique
Panic attacks or phobias	✓ Use at onset or after trigger
OCD / intrusive thoughts	✓ Paired with thought interruption
Emotional flooding	✓ Gentle reset
Children with anxiety	✓ In game form ("eye tag" or "follow the feather")
Autistic shutdown or sensory overload	✓ With adapted pacing
Freeze response / immobility	✓ With grounding or rhythmic touch

References

- Shapiro F. Eye Movement Desensitization and Reprocessing: Basic Principles, Protocols, and Procedures. Guilford Press; 2001.
- Stickgold R. EMDR: A putative neurobiological mechanism of action. J Clin Psychol. 2002;58(1):61-75.
- Propper RE, Christman SD. Interhemispheric interaction and saccadic horizontal eye movements. Brain Cogn. 2008;68(2):84-90.
- Porges SW. The Pocket Guide to the Polyvagal Theory. Norton; 2017.

Upward Eye Movements: Lifting Prefrontal Access

“When the eyes rise, the brain follows — from shutdown to spark.”

Scientific Basis

Upward gaze activates key brain regions involved in alertness, executive function, and re-engagement:

- Midline prefrontal cortex – supports planning, decision-making, and working memory.
- Reticular activating system (RAS) – engaged via the superior colliculi to increase wakefulness.
- Cortical-brainstem loops – linked to orienting responses and re-entry from freeze states.

When the body drops into dorsal vagal shutdown or depressive immobility, clients often:

- Gaze downward or fixate without blinking.
- Lose access to verbal and cognitive processing.
- Appear flat, disconnected, or “checked out.”

Gently lifting the gaze upward interrupts this pattern, helping restore cognitive access, awareness, and motor readiness by “waking up” suppressed circuits.

Clinical Purpose

Upward eye movements are a gentle activation tool to:

- Re-engage executive brain regions in freeze, fatigue, or collapse.
- Interrupt immobilization and early dissociation.
- Stimulate top-down access for verbal memory or goal-directed action.
- Increase alertness in hypoaroused clients.
- Support a transition from internal withdrawal to external orientation.

How to Do It

Guided Protocol: Upward Gaze Activation

1. Invite the client to sit comfortably, ideally with back support.
2. Ask them to lift only their eyes (not their head) toward the upper third of the visual field.
3. Hold gaze for 3–5 seconds, then return to neutral.
4. Repeat for 5–7 gentle lifts, pausing between each one.
5. Add a breath cue: “As you look up, take a soft inhale. As you return, exhale slowly.”

Variations

- Combine with cross-lateral movement (e.g., tapping opposite knee or shoulder while eyes lift).
- Use a target dot or small object placed just above eye level as a visual cue.

Therapist Tip: Upward gaze should always feel gentle and non-straining. If a client feels eye fatigue, dizziness, or discomfort, shorten holds or use fewer repetitions.

How to use

Use Case	Recommendation
During freeze/dissociation episodes	3-5 lifts, spaced with grounding touch
To exit shutdown or fatigue	1-2 rounds upon waking or during therapy

Practical Ways to Integrate into Everyday Life

- Morning wake-up: Before getting out of bed, lift your eyes softly 3-5 times to “switch on” the brain for the day.
- Desk or study breaks: Look up toward the ceiling or a window frame every hour to bring alertness back online.
- While brushing teeth or standing in the kitchen: Use a point above eye level (a shelf, cabinet edge) for a few lifts to gently re-engage focus.
- Transition moments: Before starting emails, therapy notes, or planning sessions, do a few lifts to help the prefrontal cortex “come online.”
- Nature moments: Watch a bird flying overhead, a tree swaying, or clouds drifting high — letting the gaze rise without strain.

Observable Signs of Regulation

- Blinking resumes
- Verbal processing improves
- Increased tone or posture shift ("coming back online")
- Color returns to face
- Return of emotional access ("I can feel something again")
- Ability to engage in therapy resumes after collapse

Best Matches Per Diagnosis / Symptom Cluster

Diagnosis / Pattern	Suitability
Chronic fatigue or burnout	✓ Gentle pre-activation
Chronic fatigue or burnout	✓ Gentle pre-activation
Grief / immobilizing sadness	✓ With safe holding
ADHD hypoactive subtype	✓ Alerting without overstimulation
Executive dysfunction / trauma shutdown	✓ Bridges prefrontal access
Children who 'zone out'	✓ In play form ("spot the ceiling fairy")

References

- Lanius R, et al. The neurobiology of trauma and dissociation: Towards an integrative understanding. Eur J Psychotraumatol. 2018;9(1):1472981.
- Ogden P, Fisher J. Sensorimotor Psychotherapy: Interventions for Trauma and Attachment. Norton; 2015.
- Schore AN. Affect Regulation and the Repair of the Self. Norton; 2003.
- Porges SW. The Polyvagal Theory. Norton; 2011.

Converging Gaze: Midline Integration and Vagal Tone Activation

"When the eyes meet at center, the nervous system remembers itself."

Scientific Basis

Converging gaze — focusing both eyes on a single, close midline target — stimulates multiple pathways that support parasympathetic activation and integration:

- Oculomotor nerve (cranial nerve III) – shares brainstem origins with vagal nuclei, linking eye position to vagal tone.
- Anterior cingulate cortex & insula – support emotion regulation, attention, and interoceptive awareness.
- Ventral vagus nerve – gently activated when convergence is combined with slow nasal breathing.
- Corpus callosum – midline-focused eye work promotes left-right hemispheric balance.

This form of eye positioning has been used in yogic *trataka*, convergence exercises in neuro-optometry, and modern vagal regulation practices.

Clinical Purpose

Converging gaze is a centering tool that can:

- Anchor ventral vagal tone after stress or activation.
- Support midline integration (balancing left and right hemispheres).
- Shift clients from chaos or shutdown into centered calm.
- Enhance interoceptive awareness (connection to body signals).
- Help regulate visual instability in trauma or sensory processing disorders.

How to Do It

Midline Convergence Protocol

1. Ask the client to hold a pen or thumb about 15–20 cm (6–8 inches) in front of their nose.
2. Invite them to softly gaze at the object — no straining or forcing focus.
3. Hold gaze for 10–20 seconds, then relax the eyes.
4. Repeat for 3–5 cycles, encouraging slow nasal breathing.
5. Option: layer with humming, a hand on the chest, or light tapping for vagal co-activation.

Therapist Tip: Cue gently — "Let your eyes meet at center, no force, just soft contact." Eyes may tremble or shift; that's normal. Simply guide them back to center.

How to use for regulation

Use Case	Recommendation
Daily vagal tone support	1-2 rounds morning and evening
During therapy closure	Anchor safety before session ends

Practical Ways to Integrate into Everyday Life

- Morning center: Hold a coffee mug or toothbrush at nose level and let your eyes softly meet at center for a few seconds.
- Desk breaks: Focus on a pen or finger at midline between emails to settle your system.
- Nature moments: Bring a flower, leaf, or pebble close and focus softly on it.
- Transition rituals: Before meditation, journaling, or prayer, use 2-3 cycles to “come into yourself.”
- Evening grounding: Use a candle or soft light at midline for a short gaze practice to calm before bed.

Observable Signs of Regulation

- Breath slows naturally
- Increased salivation (sign of vagal tone)
- Gentle swaying or grounding sensations
- Eye blinking normalizes
- Return to emotional safety (“I feel centered”)
- Warmth or tingling in chest and core

Best Matches Per Diagnosis / Symptom Cluster

Diagnosis / Pattern	Suitability
Anxiety with hyper-visual scanning	✔ Shifts to inward focus
Sensory processing disorders	✔ Supports visual recalibration
Autonomic dysregulation (POTS, IBS)	✔ Engages vagal reflex loops
ADHD or attention fragmentation	✔ Helps build sustained visual focus
Shutdown or floating states	✔ Anchors into body midline awareness

References

- Lehrer P, Gevirtz R. Heart rate variability biofeedback: how and why it works. Front Psychol. 2014;5:756.
- Balaban CD, Thayer JF. Neurological bases for balance–anxiety links. J Anxiety Disord. 2001;15(1-2):53-79.
- Porges SW. The Polyvagal Theory. Norton; 2011.
- Simons JP, et al. Visual fixation and vagal tone modulation: insights from meditation and gaze studies. J Psychophysiol. 2019;33(3):201-213.

Still-Point Gaze Calming the System Through Eye Stillness

“Stillness in the eyes creates stillness in the body. From overwhelm to presence, one gaze at a time.”

Scientific Basis

When the eyes stop moving and softly hold on one point, the nervous system receives a powerful cue to settle.

- The vestibulo-ocular reflex quiets, creating a sense of balance.
- The reticular activating system (RAS) downshifts, reducing arousal.
- The limbic system eases its alert output.

This combination:

- Lowers neural excitation in the brainstem and thalamus
- Increases parasympathetic activity by reducing constant “micro-scanning”
- Supports the alpha/theta brainwave states linked to calm and presence

This practice is seen across traditions — from yoga drishti to craniosacral still points — as a way to guide the body from alertness into quiet safety.

Clinical Purpose

Still-point gaze can:

- Calm an overstimulated nervous system without adding more input
- Help clients in freeze or collapse states reconnect to body awareness
- Create a bottom-up anchor for moments of overwhelm
- Offer a sense of containment and safety when words aren’t possible

How to Do It

Still-Point Protocol

1. Invite the client to choose a single spot (dot on wall, candle flame, grain in wood).
2. Ask them to rest their eyes on that point — no scanning, no darting.
3. Encourage soft stillness (no forced blinking, no strain).
4. Hold gaze for 20–60 seconds, pairing with slow breath.
5. Ask them to notice sensations (weight, grounding, quietness).

Cue: “Let your eyes find one point to rest on — like your gaze is settling into a soft nest.”

Duration and Frequency

Use Case	Recommendation
During emotional overwhelm	30- seconds, repeat up to 3 rounds
Daily regulation support	2-3 minutes of still gaze meditation
Before or after triggering discussion	Short pause with silent gaze hold

Practical Ways to Integrate into Everyday Life

- Morning moment: Rest your gaze on a single point in the room (e.g., a picture frame corner) for 30 seconds before starting your day.
- Coffee or tea break: Soften your eyes onto the steam rising from your mug or a single spot on the table to let your system settle.
- At your desk: Pick a fixed point – a pin on a corkboard, a small object – and rest your gaze there when you feel mentally cluttered.
- Nature pause: Fix your eyes on a flower, leaf, or stone on the ground, allowing the stillness to slow your breath and thoughts.
- Before bed: Focus softly on a candle flame, the edge of a shelf, or a point on the wall for 30-60 seconds to cue the body into quiet before sleep.

Observable Signs of Regulation

- Subtle bodily stillness (less fidgeting)
- Lowered shoulders or deep sigh
- Eyes soften or moisten
- Facial tone becomes neutral or open
- Emergence of emotion without overwhelm
- Client reports feeling "quieter," "heavier," or "more here"

Best Matches Per Diagnosis / Symptom Cluster

Diagnosis / Pattern	Suitability
Dorsal vagal shutdown / freeze	✓ First-line tool
Autonomic collapse or burnout	✓ Non-effortful entry point
Panic post-collapse stage	✓ Re-entry to felt sense
High sensitivity / neurodivergence	✓ Safe visual anchor
Clients with poor body awareness	✓ Restores interoceptive felt sense

References

- Ogden P, Minton K, Pain C. Trauma and the Body: A Sensorimotor Approach to Psychotherapy. Norton; 2006.
- Scaer RC. The Trauma Spectrum: Hidden Wounds and Human Resiliency. Norton; 2005.
- Porges SW. Clinical Applications of the Polyvagal Theory. Norton; 2018.
- Siegel DJ. The Mindful Brain. Norton; 2007.

Somatic Regulation

"The body remembers — and the body restores."

These tools are not about exercise or fitness — they are small, intentional movements and sensations designed to shift nervous system state safely.

Reflection: Soma – The Living Body Within

Soma.

From the ancient Greek word for body — but not just any body. Soma is the living, felt, conscious body: the body as experienced from the inside out, the landscape of emotions, sensations, intuition, and memory that is uniquely yours.

There's a saying: "Thoughts are the language of the brain, and emotions are the language of the body."

Somatic work is learning to listen to that language again — to reconnect with the body not as an object to be managed, but as a living home to be re-inhabited.

Soma is where trauma lives — not as stories, but as unfinished movements, frozen patterns, and stored tension. The jaw that never unclenched, the breath that never fully left, the shoulders still braced for impact.

And yet, soma is also where healing begins.

When we return with safety, rhythm, and presence, the body softens. The tissues reawaken. The held breath releases.

As Peter Levine reminds us: "Trauma is in the nervous system, not the event."

In Greek philosophy, soma was seen as the vessel of the soul — the temple through which your inner world takes form in the outer one.

Somatic awareness is a door into interoception — the deep noticing of what your body is whispering beneath words. It's a key to regulating the autonomic nervous system, restoring felt safety, and raising vagal tone.

Soma is not an object. It is not a diagnosis.

It is the felt, breathing, sensing, knowing body — the place where your stories are stored, your wounds are held, and your healing waits.

Introduction to Somatic Regulation

Why Somatic Regulation Matters

1. Somatic regulation refers to therapeutic techniques that engage the body directly — through movement, touch, posture, or sensation — to influence the autonomic nervous system (ANS) and restore safety, presence, and integration.

Unlike cognitive or verbal approaches, somatic tools:

1. Bypass the thinking brain (prefrontal cortex)
2. Communicate directly with brainstem and limbic structures
3. Activate the vagus nerve and interoceptive pathways
4. Help resolve trauma stored in procedural memory and sensory systems (1)

Somatic regulation is essential in trauma-informed care because many clients:

1. Cannot access emotions through words
2. Experience shutdown or freeze in talk-based therapy
3. Need bottom-up safety cues to feel grounded and connected

What Makes a Regulation Practice Somatic?

A technique is somatic when it:

1. Involves movement, touch, or sensation awareness
2. Works through the body's natural rhythms (e.g., swaying, rocking, pressure)
3. Affects neurochemical states (e.g., cortisol, oxytocin, dopamine)
4. Builds capacity in the Window of Tolerance through embodied repetition (2)
5. Doesn't require storytelling or cognitive insight to be effective

Somatic Tools in Nervous System Work

In this module, you'll learn how to use tools such as:

1. Grounding sequences (e.g., pressing feet, wall push)
2. Proprioceptive input (e.g., pressure, joint compression)
3. Vestibular input (e.g., swaying, head tilts, rocking)
4. Tactile regulation (e.g., self-hold, hand-over-heart, textured tools)
5. Bilateral movement (e.g., tapping, walking, cross-crawl)
6. Each technique is paired with:
7. Clear clinical indications
8. Contraindications and safety considerations
9. Diagnostic matches
10. Client-friendly language to guide use

Scientific Foundations

1. Polyvagal theory shows how movement and body signals shift vagal tone and safety perception (3)
2. Somatosensory cortex activation (through pressure/touch) helps anchor presence
3. Brainstem-cerebellar circuits modulate arousal via posture, balance, and motor feedback (4)
4. Mirror neurons and co-regulation reinforce embodied healing through modeled movements

When to Use Somatic Regulation

Scenario	Somatic Entry Point
Client is “in their head”	Body-based grounding resets interoception
Talk therapy is stuck	Movement unlocks stored trauma loops
Emotional overwhelm	Proprioceptive or tactile input calms limbic reactivity
Panic or dissociation	Simple sensory anchors return awareness
Chronic pain or IBS	Bottom-up calming shifts vagal tone and reduces threat signals

References

- van der Kolk B. The Body Keeps the Score. Penguin; 2014.
- Levine PA. Waking the Tiger: Healing Trauma. North Atlantic Books; 1997.
- Porges SW. The Pocket Guide to the Polyvagal Theory. Norton; 2017.
- Schore AN. Affect Regulation and the Repair of the Self. Norton; 2003.
- Ogden P, Minton K, Pain C. Trauma and the Body. Norton; 2006.

Grounding Techniques: Returning to the Body Through Contact and Gravity

"Before we can rise, we must first arrive. Grounding is the body's way of saying, 'I am here.'"

Scientific Basis

Grounding techniques involve intentional contact with the earth, environment, or the body's own weight to stabilize the nervous system. These inputs:

- Stimulate proprioceptive and interoceptive systems, enhancing body awareness (1)
- Activate pressure receptors and slow-adapting tactile neurons, which feed into the vagal brake (2)
- Regulate brainstem arousal and increase insula and anterior cingulate activity, promoting emotional presence (3)
- Re-establish the body's relationship with gravity, an anchor often lost in dissociation or sympathetic overdrive

Clinical Purpose

- Restore presence during emotional overwhelm, panic, or freeze
- Anchor the client in 'felt sense' safety through pressure and contact
- Help integrate bottom-up and top-down inputs
- Support transitions into or out of breath, eye, or cognitive-based work
- Improve body schema in clients with trauma, sensory sensitivity, or dissociation

Reflection: Grounding - The Anchor of Being

Trauma pulls us up and out. Grounding brings us down and in.

To feel your feet on the earth — to notice weight, pressure, gravity — is to return to your body after drifting into your head, or hovering outside yourself for too long.

In Hebrew, the root word for ground is also the root of the word human.

To ground is to remember that you belong here.

Research shows that grounding can calm inflammation, cortisol, and even improve sleep. But beyond the science, it does something simpler and more immediate: it gives you an anchor.

And an anchor changes everything.

When the storm hits, an anchor keeps you from floating away. When the waves crash, you have something to lean against. Grounding announces you — it's a kind of super pose.

It says: I'm here. I'm backing myself.

Grounding also carries a symbolic truth. It represents stability — to be unshaken. It is rooting — to connect yourself back to your own story. It is belonging — being at home in your own body.

And it is humility — for in Hebrew, humility shares its root with earth. Grounding restores the right relationship between myself and creation.

Grounding doesn't just steady the body — it speaks to the brain. It re-engages interoception, proprioception, and neuroception — the senses that remind your nervous system that you are safe, present, and whole.

To ground is not just to stand on the earth. It is to stand in yourself.

How to Do It:

Foundational Grounding Exercises

Feet on the Floor

Ask the client to press their bare or socked feet into the floor.

Invite awareness of the weight, texture, and support.

Cue: "Let your feet remind you where you are."

Wall Press or Door Frame Lean

The client stands facing a wall or inside a doorway, gently pushing hands into it.

Hold pressure for 5-10 seconds, then release slowly.

Repeat 3-5 rounds.

Cue: "Feel the power of your body meeting the world."

Sit and Drop

Client sits on a firm surface and gently drops awareness into their hips and sit bones.

Allow breath to follow.

Optional: hold small weighted object (like a sandbag or cushion) on lap.

Grounding Scan

Move attention through anchor points (feet, hips, back, hands).

Combine with slow breath.

Cue: "Where do you feel most connected right now?"

Duration and Frequency

Context	Recommendation
During panic or dissociation	Immediate 1-2 minute intervention
Start of therapy or session	3-5 minutes to enter window of tolerance
Daily self-regulation	Morning/bedtime grounding for nervous system tone

Contraindications and Cautions

Concern	Note
Hypervigilance or trauma linked to body contact	Offer guided visual grounding or less direct pressure
Vestibular dysregulation	Avoid positions that shift head quickly or reduce stability
Tactile defensiveness	Use environmental anchors (chair, surface) over self-touch

Observable Signs of Regulation

- 1.Shoulders drop, breath slows
- 2.Client begins to orient to present moment
- 3.Muscle tension reduces
- 4.Return of emotion or tears (especially in freeze states)
- 5.Feet, hands, or torso become warmer (“I feel back in my body”)
- 6.Facial expression softens

Best Matches Per Diagnosis / Symptom Cluster

Diagnosis / Pattern	Suitability
Panic attacks / anxiety	✔ Anchor out of hyperarousal
Dissociation / freeze	✔ Safe re-entry to body
Sensory integration challenges	✔ Body schema recalibration
Trauma flashbacks	✔ Disruption of disembodied narrative
Children or non-verbal clients	✔ Universally accessible regulation tool

References

- Mehling WE, et al. Body awareness: a phenomenological inquiry. J Altern Complement Med. 2011;17(5):349-356.
- McGlone F, et al. Discriminative and affective touch: Sensing and feeling. Neuron. 2014;82(4):737-755.
- Craig AD. How do you feel? Interoception: the sense of the physiological condition of the body. Nat Rev Neurosci. 2002;3(8):655-666.
- Porges SW. The Pocket Guide to the Polyvagal Theory. Norton; 2017.
- Levine PA. In an Unspoken Voice. North Atlantic Books; 2010.

Proprioceptive Regulation: Pressure, Resistance, and Body Integrity

"Through resistance, we reclaim presence. Pressure is not the enemy of calm — it is its architect."

Scientific Basis

Proprioception is the body's internal sense of position, movement, and resistance. It's mediated by muscle spindles, joint capsules, and Golgi tendon organs, which send information to the cerebellum, brainstem, and somatosensory cortex (1). Proprioceptive input is:

- Organizing and calming, especially for children or clients in sensory overload
- A powerful vagal stimulator when delivered through sustained, rhythmic pressure
- Central to re-establishing body boundaries in trauma recovery
- A key regulator for emotional self-regulation and interoceptive accuracy (2,3)

In trauma or dysregulation, proprioception is often dulled or fragmented, leaving clients disconnected from their felt sense. Proprioceptive tools rebuild internal coherence.

Clinical Purpose

- Reconnect body map (somatosensory cortex) and improve interoceptive accuracy
- Calm sympathetic arousal or sensory chaos
- Establish safe body boundaries in clients with trauma
- Improve tone, posture, and body orientation in physical and emotional dysregulation
- Build regulation capacity in children and neurodiverse individuals

How to Do It

Proprioceptive Input Techniques

Weighted Blanket or Compression Garments

- 5-10% of body weight for blankets; compressive socks, vests, or sleeves for subtle input
- Use during seated tasks, sleep, or regulation routines

Wall Pushes

- Client stands arms-length from wall, leans and presses with palms (5-10 seconds)
- Release and feel body weight; repeat 3-5 times
- Variations: push through legs or shoulders

Chair Pull or Arm Press

- Sitting in a firm chair, client presses arms into armrests or pulls up on seat edge
- Focus on slow, steady pressure

Therapist-Guided Resistance

- Hold a resistance band or soft object between hands and gently pull apart
- Combine with breathing or eye focus for layered regulation

Animal Walks / Crawling (for kids)

Bear walks, crab crawls, or pushing weighted objects builds full-body input

Duration and Frequency

Context	Recommendation
Acute dysregulation	30 seconds to 2 minutes, repeat as needed
Therapeutic pairing	Before or after emotional processing work
Daily self-regulation	5-10 minutes total in routines

Contraindications and Cautions

Concern	Note
Joint hypermobility	Modify techniques; use compression rather than stretch
Chronic pain conditions	Avoid deep pressure unless clearly tolerated
Aggressive dysregulation	Avoid stimulating resistive play until grounded

Reflection: Proprioception – How the Soul Finds Its Temple

Proprioception gives walls to words and emotions.

It is the unsung sixth sense – the one that tells you where you are in space without needing to look. It comes from the Latin word proprio, meaning one's own.

Proprioception is how your brain senses position, movement, tension – the gentle pull in a tendon, the quiet stretch in a joint – all without sight. It is the subtle conversation between muscles and mind that whispers: "This is where you are."

Trauma interrupts that whisper. It makes you feel unmoored, floating, untethered.

Proprioception restores the borders. It provides feedback, boundaries, and safety. When your nervous system feels where the edges are, it can finally rest inside them.

Proprioception is how the soul locates its temple.

You don't just think it – you know it in your bones.

Observable Signs of Regulation

1. Breathing slows or deepens
2. Shoulders relax, posture becomes upright
3. Verbal coherence improves
4. Client reports feeling “contained,” “held,” “stronger,” or “calmer”
5. Reduction in chaotic movements or emotional reactivity
6. Children show increased focus and play engagement

Best Matches Per Diagnosis / Symptom Cluster

Diagnosis / Pattern	Suitability
ADHD / sensory modulation disorders	✓ Strong regulator of tone and attention
Autism spectrum conditions	✓ Organizes sensory chaos when tailored
Trauma with disembodiment	✓ Anchors body map and integrity
Panic / hyperarousal	✓ Lowers over-excitation safely
Children with melt-downs	✓ Quickly restores control and connection

References

- Schaaf RC, Benevides T, Mailloux Z. Proprioception: A Multisystem Process. *Front Integr Neurosci.* 2015;9:32.
- Dunn W. The Sensory Profile: Measuring patterns of sensory processing. *Clin Percept Mot Skills.* 1999;89(3):774-786.
- Porges SW. *The Pocket Guide to the Polyvagal Theory.* Norton; 2017.
- Fisher J. *Sensorimotor Psychotherapy.* Norton; 2015.
- Siegel DJ. *The Developing Mind.* Norton; 2012.

Vestibular Regulation: Swaying, Rocking, and Rebalancing the Inner World

“The body’s rhythm is a lullaby for the brain.”

Scientific Basis

The vestibular system, housed in the inner ear, helps regulate balance, orientation, and movement in space. It is intricately connected to:

- The autonomic nervous system (ANS) — especially arousal, orientation, and vagal tone
- Cerebellar and brainstem nuclei, responsible for regulation of posture, movement, and arousal thresholds
- Emotional centers including the amygdala and hippocampus (1)

Vestibular input (e.g., swaying, spinning, rocking) is one of the earliest sensory experiences in human development — think of a baby being rocked to sleep. It profoundly impacts:

- State regulation
- Safety perception
- Emotion modulation
- Social engagement readiness (2,3)

Clinical Purpose

- Reset disrupted arousal levels in trauma and sensory dysregulation
- Support self-soothing through rhythmic movement
- Facilitate transitions between nervous system states (e.g., from high alert to calm)
- Re-establish a body-in-space awareness, often lost in trauma or dissociation
- Improve focus, balance, and regulation capacity in children and adults alike

How to Do It

Vestibular Regulation Techniques

Seated Rocking

- Sit on a firm chair or therapy ball. Gently rock forward-back or side-to-side.
- Cue: “Let your body find its own rhythm — like waves in the ocean.”
- Ideal before or after intense emotional work.

Slow Head Turns

- Instruct client to gently turn head left to right with breath.
- Match to slow inhale/exhale. Can be paired with eye movement tracking.

Swaying While Standing

- Feet hip-width apart. Let body gently sway in figure 8 or lateral rhythm.
- Eyes open or softly closed.

Hammocks / Therapy Swings (for children or adults in sensory rooms)

1. Gentle, linear motion preferred over spinning.
2. Always supervised and kept slow, rhythmic, and predictable.

Partner Sway

- Therapist or caregiver can support a light sway movement, creating co-regulation.
- Cue: “Let’s breathe and move together — slowly and safely.”

Duration and Frequency

Context	Recommendation
Daily nervous system hygiene	2-3 minutes swaying/rocking
During dysregulation	Immediate use for calming or alerting
Pre-session	1-2 minutes to prime attention and body awareness

Contraindications and Cautions

Concern	Note
Vertigo, vestibular disorders, or motion sickness	Avoid or limit input; consult physician
Post-concussion syndrome	Vestibular input may worsen symptoms
PTSD with dizziness triggers	Use minimal, linear input only
Hyperactivity or over-arousal	Fast spinning or irregular movement may overstimulate – keep slow and rhythmic

Reflection: Vestibular – The Body's Inner Compass

When the world tilts, the body remembers gravity.

When the mind spirals, the inner ear listens for center.

The vestibular system – deep in the inner ear – is your hidden compass, always asking, “Where am I in space? Am I upright, moving, falling, or still?”

To reorient is not just to stand – it is to return to knowing where you are. It is a quiet reunion between body and earth.

The vestibular system is the lullaby for the brain. The sway of a hammock, the rocking of a cradle, the gentle turn of a dance – they soothe not only the body, but the mind. We are wired from birth to calm when we are rhythmically moved.

Trauma can disrupt this harmony. It leaves us dizzy, disoriented, unsure of our footing – like the inner compass has lost its true north.

But when you reawaken the vestibular sense – with slow turns, gentle sways, and intentional movement – the world steadies. The brain exhales. You remember center.

And in that moment, balance is no longer just physical – it becomes a way to come home to yourself.

Observable Signs of Regulation

- 1.Eyes soften or close naturally
- 2.Breathing synchronizes with movement
- 3.Increased yawning, sighing, or tearfulness (release)
- 4.Smiling, grounded posture, or emotional accessibility returns
- 5.Improved orientation to present time/place

Best Matches Per Diagnosis / Symptom Cluster

Diagnosis / Pattern	Suitability
ADHD / attention dysregulation	✔ Builds rhythm and focus
PTSD / trauma loops	✔ Resets arousal and helps reintegrate body map
Shutdown, freeze, or dissociation	✔ Gently restores movement and presence
Anxious restlessness	✔ Rhythmic input soothes overactivation

References

- Shaffer F, McCraty R, Zerr CL. A healthy heart is not a metronome: an integrative review of HRV. Front Psychol. 2014;5:1040.
- Ayres AJ. Sensory Integration and the Child. Western Psychological Services; 2005.
- Levine PA. In an Unspoken Voice. North Atlantic Books; 2010.
- Blakeslee S, Blakeslee M. The Body Has a Mind of Its Own. Random House; 2007.
- Porges SW. The Polyvagal Theory. Norton; 2011.

Tactile Regulation: Anchoring Through Touch, Texture, and Containment

"Safe touch tells the body: you exist, you're real, and you're not alone."

Scientific Basis

Tactile regulation uses intentional, safe forms of touch and texture-based sensory input to calm the nervous system. This works by stimulating:

- C-tactile (CT) afferent fibers, which are activated by slow, gentle touch and linked to emotional soothing (1)
- The insular cortex, which integrates interoception, emotion, and body-based safety perception (2)
- Oxytocin release, enhancing social bonding, pain modulation, and stress reduction (3)
- Somatosensory cortex and vagal pathways, helping re-anchor the body in real-time space

Touch-based techniques support co-regulation, emotional expression, and reconnection to a felt sense of boundary and safety – especially important in trauma recovery and early developmental repair.

Clinical Purpose

- Provide grounding and containment in dysregulated or overwhelmed states
- Restore safe body boundaries in trauma, shutdown, or fragmentation
- Co-regulate with caregiver or therapist through touch
- Activate oxytocin and parasympathetic tone through gentle, rhythmic tactile input
- Build interoceptive awareness when cognitive or verbal processing is inaccessible

How to Do It

Tactile Regulation Techniques

Self-Containment Hold

- Client gently places one hand on chest, one on belly.
- Inhale: feel hands rise. Exhale: soften shoulders.
- Invite awareness of breath, heartbeat, or warmth.
- Cue: "Your own touch can remind you that you're here."

Hand Over Heart Technique

- One hand over sternum, soft pressure. Use during stress, anxiety, or grief.
- Optional paired mantra: "I am safe. I am home."

Textured Object Anchors

- Keep small textured objects (fabric, smooth stones, rough rubber) in pocket or on lap.
- Use in sessions, classrooms, or transitions.
- Especially useful for autism, ADHD, or anxiety.

Weighted Touch

- Therapist or caregiver applies slow, gentle pressure with a weighted cushion or hands (with consent).
- Areas: shoulders, upper back, or thighs. Duration: 30–60 seconds.
- Always ask permission and avoid trauma-linked areas.

Tapping or Brushing

Use soft brush or fingers to tap lightly down the arms, legs, or back.

Create rhythm; combine with breath or humming for multisensory calming.

Duration and Frequency

Context	Recommendation
In-session support	1-3 minutes, paired with breath or emotion work
Daily self-regulation	Morning/bedtime containment practice
Acute anxiety or collapse	Immediate tactile anchor, even for 30 seconds

Contraindications and Cautions

Concern	Note
Touch aversion / trauma history	Always offer opt-in tactile work; avoid assumptions
Tactile defensiveness	Begin with indirect or textured objects before body contact
Abuse history	Use self-touch or sensory tools, never therapist-initiated contact without prior consent
Overstimulated children	Tactile tools may need to be very soft, brief, or introduced during calm moments

Observable Signs of Regulation

1. Breath slows, shoulders drop
2. Client begins to cry or soften
3. Increased ability to speak about emotion
4. Less fidgeting or collapse; more presence and orientation
5. Child returns to task or rest posture
6. Client reports: "That felt comforting," "I feel more solid," "I needed that."

Best Matches Per Diagnosis / Symptom Cluster

Diagnosis / Pattern	Suitability
Complex trauma	✓ With resourcing and body consent
Anxiety / panic	✓ Safe self-hold or object anchoring
Dissociation	✓ Self-touch or texture to re-enter body
Autism / sensory modulation	✓ With personalized object use
Children in distress	✓ Touch + weight often stabilizes emotion

Reflection: The Language of Touch

Touch is the first sense we ever know.

Before we see or hear, we are felt — cradled, held, wrapped in warmth. The skin becomes our first teacher, telling us if the world is safe, if love is present, if we belong.

The tactile system is not just about pressure and texture — it is the nervous system's oldest form of conversation.

Every stroke, every press, every soft brush of fabric or fingertip speaks directly to the brain.

Trauma can make this conversation confusing. For some, touch becomes too loud — the slightest contact feels like an alarm. For others, touch becomes numb, as if the body stopped listening altogether.

Yet touch has the power to regulate and restore.

- *Deep, steady pressure can slow the heart rate, quiet the amygdala, and release oxytocin — the hormone of trust and safety.*
- *Gentle, rhythmic touch (like hand-on-heart, a weighted blanket, or a slow self-hold) can anchor the body, reminding the nervous system that it is safe to soften.*

There's a beautiful paradox in touch: it sets boundaries and dissolves barriers. It says, "This is where you end and the world begins," while also saying, "You are not alone."

The tactile system whispers to every cell: "You are here. You are held."

References

- McGlone F, Wessberg J, Olausson H. Discriminative and affective touch. *Neuron*. 2014;82(4):737-755.
- Craig AD. Interoception: the sense of the physiological condition of the body. *Curr Opin Neurobiol*. 2003;13(4):500-505.
- Uvnäs-Moberg K, Petersson M. The neuroendocrinology of oxytocin. *Brain Res*. 2005;36(2):225-233.
- Ogden P, Fisher J. *Sensorimotor Psychotherapy*. Norton; 2015.
- Porges SW. *Clinical Applications of the Polyvagal Theory*. Norton; 2018.

Bilateral & Rhythmic Regulation: Cross-Body Integration for Trauma and Focus

“Trauma fragments. Rhythm restores. Bilateral movement weaves the pieces back together.”

Scientific Basis

Bilateral and rhythmic regulation uses alternating left-right movement, stimulation, or tapping to:

- Engage both hemispheres of the brain, supporting integration of trauma memories and sensory input (1)
- Stimulate the corpus callosum, which facilitates interhemispheric communication
- Reorganize sensorimotor loops in the body and brain
- Support coherence in the reticular activating system, improving focus and regulation (2)
- Mimic natural regulatory rhythms (e.g., walking, crawling, rocking), which evolved to calm the nervous system (3)

Bilateral input is also foundational to therapies like EMDR, where rhythmic eye or tactile stimulation helps reprocess traumatic material in a regulated state.

Clinical Purpose

- Re-integrate fragmented sensory/emotional experiences
- Calm emotional flooding or hyperactivation
- Improve focus, sequencing, and body awareness in ADHD, trauma, and learning disorders
- Activate cross-lateral brain engagement in neurodevelopmental support
- Help access verbalization, emotional clarity, and embodied presence during therapy

How to Do It

Bilateral Regulation Techniques

Alternating Tapping

- Client taps left and right sides of body (knees, shoulders) in rhythmic pattern
- Can be self-administered or therapist-guided
- Use steady rhythm: ~1 tap per second
- Optional verbal pairing: “Safe / Here” or “Left / Right”

Cross-Crawl Marching

- While standing or seated, touch left hand to right knee and vice versa
- Match to slow breath
- Use before focus tasks, school transitions, or therapeutic re-engagement

Butterfly Hug

- Cross arms over chest, fingers on collarbones or upper arms
- Tap alternately with gentle pressure
- Cue: “This is your body remembering safety.”

Rhythmic Object Passing

- Pass a weighted object left to right across midline
- Combine with counting, music, or storytelling for cognitive-emotional integration

Walking or Drumming

Walking in nature, drumming beats, or dancing with bilateral rhythm supports both nervous system tone and emotional rhythm

Duration and Frequency

Context	Recommendation
In-session trauma work	30 sec to 2 minutes between emotional peaks
Focus or grounding prep	1-3 minutes before learning or therapy
Home routines	Use during transitions, emotional regulation moments

Contraindications and Cautions

Concern	Note
Severe trauma or disorientation	Start slow, with simple tactile forms (e.g., tapping hands)
Seizure disorders or vestibular instability	Avoid rapid movement or intense cross-body input
Autistic shutdown	May need gentle, structured input with soft cues
Sensory overload	Simplify and reduce rhythmic intensity

Reflection: The Body's Weaving Wisdom

Bilateral movement weaves the pieces back together.

There is a quiet beauty in bilateral integration — left and right, limb to limb, thought and emotion, speech and silence, logic and longing. It's like art with the body, painting unseen pictures across the midline of your life.

It's not just movement. It is a rhythm that remembers who you are.

Each cross-crawl, each mirrored motion, is like grabbing life with both hands — turning fragments into flow, turning knowledge into wisdom. Crawling, walking, swaying, drumming — these simple, primal rhythms bring you back into the pattern of your own becoming.

Bilateral rhythm does what trauma disrupts: it fuses the hemispheres, reconnects left and right brain, and integrates what has been split by survival. It reweaves what was torn.

There is something sacred in this repetition — like surrendering to the regenerative intelligence of the body. You don't have to figure it out. You don't have to solve the story. This work is subcortical — moving beneath understanding, like an aspirin dissolving into water. It simply knows what to do.

Bilateral movement becomes a language that can contain words that once felt too heavy, transforming them into action, presence, and rhythm. It can rewrite the story that trauma etched into the tissues.

Every trauma in the body dims the body's light — its vitality, its energy. And yet science tells us that a cubic centimeter of mitochondria shines brighter than a cubic centimeter of the sun. Rhythm coaxes that light back, restoring energy, coherence, and life.

And still, so many of us feel shame about movement. We shrink from its awkwardness, its exposure, as though moving freely reveals too much. But here lies the key: bilateral movement speaks to shame. It melts it, rewires it, invites safety to replace it.

Bilateral rhythm doesn't demand you understand. It simply moves the pieces back into place. And when it does, the body stops bracing and starts becoming whole again.

This is the body turning toward itself. This is rhythm as medicine.

Observable Signs of Regulation

1. Improved speech, clarity, and verbal access
2. Shoulders drop, breath deepens
3. Emotional release (e.g., sigh, tears) followed by calm
4. Eyes become focused or soften
5. Children return to task, stop repetitive movement, or express themselves better

Best Matches Per Diagnosis / Symptom Cluster

Diagnosis / Pattern	Suitability
PTSD / trauma fragmentation	✓ Key integrative support technique
ADHD / learning delays	✓ Enhances focus and sequencing
Emotional flooding / rage states	✓ Grounds and distributes energy
Anxiety / panic	✓ Redirects looped thoughts
Shutdown / freeze	✓ Brings movement into stillness

References

- Shapiro F. EMDR: Basic Principles, Protocols, and Procedures. Guilford Press; 2001.
- Siegel DJ. The Whole-Brain Child. Bantam; 2011.
- Perry BD. Rhythm, relationships and regulation: The three R's of trauma treatment. Child Trauma Acad; 2006.
- Ogden P, Fisher J. Sensorimotor Psychotherapy. Norton; 2015.
- van der Kolk B. The Body Keeps the Score. Penguin; 2014.

Limbic Discharge (Shake Off Reflex)

Technique:

- Stand with knees slightly bent
- Begin bouncing gently through the heels
- Let arms and shoulders shake freely
- Allow involuntary movement if it arises (yawning, sighing, spontaneous tremor)

Why:

This mimics the natural tremoring discharge seen in wild animals after survival stress (e.g., Peter Levine's observation of a gazelle trembling after escaping a predator). The process discharges excess adrenaline, norepinephrine, and cortisol, which otherwise remain trapped in the neuromuscular system.

Limbic discharge is thought to recalibrate the hypothalamic-pituitary-adrenal (HPA) axis, reduce amygdala hyperactivity, and restore parasympathetic dominance after a stress cycle.

Linked to:

- Brainstem & Periaqueductal Gray (PAG)
- Amygdala
- Hypothalamus
- Motor cortex & cerebellum

Duration: 1–3 minutes

Observe: Spontaneous yawn, sigh, tingling, heat, tear release, laughter—all signs of nervous system reset.

References

- Levine, P. A. (1997). *Waking the Tiger: Healing Trauma*. North Atlantic Books.
- Scaer, R. C. (2005). *The Trauma Spectrum: Hidden Wounds and Human Resiliency*. W. W. Norton & Company.
- van der Kolk, B. (2014). *The Body Keeps the Score*. Viking.
- Bracha, H. S. (2004). Freeze, Flight, Fight, Fright, Faint: Adaptationist perspectives on the acute stress response. *CNS Spectrums*, 9(9), 679–685.
- Payne, P., Levine, P., & Crane-Godreau, M. A. (2015). Somatic experiencing: using interoception and proprioception as core elements of trauma therapy. *Frontiers in Psychology*, 6, 93. <https://doi.org/10.3389/fpsyg.2015.00093>
- Porges, S. W. (2011). *The Polyvagal Theory: Neurophysiological Foundations of Emotions, Attachment, Communication, and Self-regulation*. Norton.

Part 8

NeuroSync Regulation Protocol flow

The NeuroSync 3R Method - Overview

STEP 1 – RECOGNIZE

Map the state before you intervene.

- Window of tolerance mapping
- HRV & BOLT scores
- Breath pattern screening
- Neuro-loop identification
- Interoceptive awareness scale
- Therapist observation (facial tone, posture, voice, eyes)

Key Clinical Question:

“Where is the system stuck right now?”

STEP 2 – RESPOND

Interrupt survival loops with bottom-up regulation.

- Coherence breathing protocols
- Extended exhale breath training
- Eye movement resets (horizontal, vertical, tracking)
- Somatic anchoring: posture, rhythm, grounding
- Co-regulation through therapist nervous system

Key Clinical Question:

“Which tool speaks to this body state right now?”

STEP 3 - RESTORE

Reinforce safety through repetition and rhythm.

- Daily self-tracking & state reflection
- Simple home practice protocols (micro-resets)
- Small daily safety anchors (breath, movement, sensory)
- Therapist modeling safety rhythm consistently

Key Clinical Question:

“How do we build flexible new pathways over time?”

Expand participation, integration, and full embodied life.

- Gradual life reintegration (participation ladders)
- Identity rebuilding through embodied success experiences
- Long-term autonomic resilience support
- Occupation-based participation models

Protocol Flow:

A clinical rhythm to map, regulate, and rewire loops

Step	Clinical Focus	Therapist's Role
1. Recognize	Identify dysregulation patterns, loop triggers, window position	Map symptoms to ANS states (e.g. freeze, fawn, collapse)
2. Reflect	Use interoception and client language to bring awareness to what's happening inside	Mirror and validate felt sense ("Something feels too much right now...")
3. Restore Rhythm	Apply breath, eye, and somatic tools to support physiological regulation	Select technique matched to nervous system state
4. Reconnect	Co-regulate through relational safety and therapist attunement	Model regulated presence, tone, posture
5. Re-enter Cognition	Invite insight, reframe, and meaning-making after regulation	Use top-down therapy tools only after bottom-up safety
6. Rewire with Repetition	Help client apply this loop-mapping outside of sessions	Encourage daily nervous system check-ins and tracking

NeuroSync Example protocol

Example Protocol

"The nervous system speaks in sensation, rhythm, and pattern—not words. This protocol restores rhythm from the bottom up."

Step 1: Grounding the Body (Proprioceptive Anchor)

Technique:

- Sit with feet flat on the floor
- Press heels gently into the ground
- Notice pressure in your legs, hips, and back

Why:

Proprioceptive input activates the dorsal column system and stimulates the parietal cortex, helping orient the body in space. Grounding interrupts dissociation and prepares the vagus nerve for parasympathetic engagement.

Linked to: Insula, Parietal cortex, Brainstem

Duration: 30–60 seconds

Step 2: Vagal Breath (0.1 Hz Coherence Breathing)

Technique:

- Inhale slowly through the nose for 5 seconds
- Exhale slowly through the nose or pursed lips for 5 seconds
- Optional: Use a visual cue (like a moving dot or app)

Why:

Breathing at this frequency entrains respiratory sinus arrhythmia and increases HRV (heart rate variability), which reflects vagal tone and flexibility. It also shifts the brain from beta to alpha rhythm and quiets sympathetic arousal.

Linked to: Vagus nerve, Heart-brain loop, Prefrontal cortex

Duration: 2–3 minutes

References: Lehrer et al. (2020); McCraty & Zayas (2014); Porges (2011)

Step 3: Bilateral Eye Tracking (Horizontal)

Technique:

- Hold a pen, object or finger 30–40 cm in front of face
- Slowly move it side to side (left ↔ right)
- Let the eyes follow without turning the head
- Or use a drawing activity – because this integrates eye movements already

Why:

This helps to down regulate the amygdala.

References: Stickgold (2002), van der Kolk (2014), Huberman Lab (2023)

Step 4: Somatic Activation (Shaking or Rhythmic Movement)

Technique Options:

- Gently bounce heels while seated or standing
- Shake arms and legs
- Sway side to side with soft knees

Why:

Rhythmic movement activates vestibular and cerebellar networks, which regulate postural tone, movement planning, and autonomic balance. Shaking also discharges accumulated cortisol and norepinephrine from motor neurons.

Linked to: Brainstem, Cerebellum, Vagal brake

Duration: 1-2 minutes

References: Levine (1997), Porges (2011), van der Kolk (2014)

Step 5: Visual Anchor (Soft Gaze + Safety Cue)

Technique:

- Choose a neutral or calming object (e.g., tree, soft light, photo)
- Gaze softly at it without effort
- Breathe naturally and notice sensation in body

Why:

The ventral vagal system is highly responsive to visual and auditory safety cues. Soft visual focus downregulates the amygdala and enhances insula activation, supporting interoception and social engagement.

Linked to: Insula, Amygdala, Visual cortex, Vagal tone

Duration: 60-90 seconds

References: Porges (2011); Damasio (2010); Stephen Porges Polyvagal Theory

Step 6: Interoceptive Check-In

Technique:

1. Ask: "What sensation is most present in my body right now?"
2. Label the sensation: pressure, tingling, warmth, stillness
3. Track it for 3 slow breaths

Why:

Interoception builds autonomic literacy. Noticing and naming internal signals strengthens connectivity in the insula, anterior cingulate cortex (ACC), and prefrontal regions—the same areas dysregulated in trauma and anxiety.

Linked to: Insula, ACC, Default Mode Network

Duration: 30-60 seconds

References: Craig (2009); Khalsa et al. (2018); Farb et al. (2013)

Closure Gesture (Integration Signal)

Technique:

- Wrap arms around yourself or place one hand on chest, one on belly
- Say silently or aloud: "I'm here. I'm safe."
- Take a final full breath in and out

Why:

This signals completion to the nervous system and activates oxytocin release, reinforcing calm, connection, and limbic safety. The self-touch activates afferent C-tactile fibers, which reinforce calm.

Linked to: Vagus nerve, Oxytocin pathways, Somatosensory cortex

Duration: 20-30 seconds

Choosing the Right Tool: A Nervous System Matching Guide

“The technique doesn’t heal – the right technique, at the right moment, in the right state, does.”

Purpose of This Guide

This matching guide helps therapists choose the most appropriate regulation tool based on a client’s presenting state, diagnosis, or observable pattern of dysregulation. It ensures interventions are state-informed, client-centered, and nervous system-aligned – not just applied generically.

Use this page:

- As a clinical quick-reference sheet
- To build your own layered intervention protocols (e.g., grounding → breath → eye movement)
- To guide session flow, daily homework, or co-regulation practices

Regulation Tool Matching Matrix

Client Presentation / Symptom Cluster	Best-Suited Regulation Tools
Panic / Anxiety / Racing thoughts	Grounding (Feet / Chair Drop)Extended Exhale Breath Lateral Eye Tracking Tactile Anchoring
Dissociation / Freeze / Collapse	Containment Hold Vestibular Sway Proprioceptive Push Upward Gaze + Diaphragmatic Breath
Emotional flooding / Rage / Irritability	Weighted Pressure Diagonal Eye Movement Bilateral Tapping Box Breath
Sensory Overload / Autism / ADHD	Textured Object Anchors Wall Press Rocking Chair / Swing Vertical Gaze
Shutdown After Burnout	Upward Gaze with Alert Breath Proprioceptive Marching Swaying with Open Eyes
Trauma / Intrusive Memories / Flashbacks	Butterfly Hug Bilateral Object Pass Coherence Breath
Grief / Sadness / Emotional Reconnection	Hand-over-Heart Touch Gentle Belly Breathing Rhythmic Movement / Walking
Children with meltdowns or overwhelm	Weighted Stuffed Animal Hold Animal Crawls Sideways Eye Tracking Sensory Brushing

Clinical Reflections: What Regulation Really Looks Like in Practice

"Healing isn't heroic. It's rhythmic, relational, and real."

Why Regulation Comes First

Before insight, before verbal processing, before emotional breakthroughs — comes safety. Nervous system regulation is the doorway to every other therapeutic success. Whether you're working with a child in meltdown, an adult stuck in shutdown, or a teen overwhelmed by racing thoughts, the capacity to shift state is the *single greatest clinical predictor of sustainable progress*.

Therapists are not just guides — we become external regulators until the client's internal rhythm returns.

Case Vignette 1 – Occupational Therapy | Adult Client with Complex Trauma

Profile:

- 42-year-old female, history of childhood trauma, CPTSD diagnosis
- Reports "not feeling in her body," sleep disruption, and emotional flooding
- Has done years of cognitive therapy with limited results

Session Focus:

- Began session with proprioceptive wall push and 3-minute sway
- Paired bilateral tapping with breathwork
- Tracked facial softening, increased presence, and spontaneous emotional release

Outcome:

Client said: "I've talked about this for years, but I finally felt it... and could let it go."

Case vignette 2 – Child Client | Sensory Processing and Anxiety

Profile:

- 8-year-old boy with sensory processing challenges, ADHD diagnosis
- Fidgets, startles easily, and struggles with transitions
- Avoidant of eye contact and easily dysregulated by noise

Regulation Plan:

- Began session in a sensory tent with compression vest
- Introduced rhythmic crawling and tactile object exploration
- Ended with eye tracking and seated weighted animal breathing

Outcome:

Child verbalized: "I feel like my engine went slow." (His own self-regulation metaphor.)

Clinical Reflections: What Regulation Really Looks Like in Practice

Case Vignette 4 – Adult Client | Panic Attack During Session

Profile:

- 40-year-old female, history of severe childhood trauma
- Diagnosed with generalized anxiety and panic disorder
- Reports frequent episodes of overwhelming anxiety and panic attacks
- Hypervigilant baseline, difficulty calming after activation
- Previous CBT and medication

In-Session Event:

- During session, client experienced an acute panic attack
- Reported rising sense of chest tightness, dizziness, and dissociative detachment
- Therapist observed rapid breathing, darting eyes, and rigid body posture

Intervention Sequence:

- Initiated lateral eye movements (horizontal tracking, left-right slow gaze shifts)
- Introduced upward right gaze, held 15 seconds
- Instructed left eye closure while maintaining right eye upward gaze (stimulating asymmetric brain activation)
- Applied cold stimulation (ice pack) to lateral neck targeting vagus nerve cooling
- Guided client into slow exhale-focused breathing (5-second exhale with brief pause)

Outcome:

- Gradual reduction in breathing rate and visible relaxation of facial tension
- Client verbalized:
- “I can feel my chest opening and I’m coming back into my body.”
- Within 4 minutes, panic symptoms resolved, client remained regulated for remainder of session

NeuroSync protocol layering:

- Eye movement → cranial nerve modulation → vagal stimulation → breath regulation → state restoration.

Closing Note to Practitioners

Every time you help a client shift from chaos to calm, from collapse to contact, from overload to rhythm — you are rewiring not just their nervous system, but their sense of agency, presence, and safety.

Start with the body. The rest will follow.

Final Reflection: The Fourth R – Remember

We learn to recognize what's happening inside, respond with the right tool, and restore balance.

But there is one more quiet invitation, a thread that ties it all together: Remember.

Remember that your nervous system is not just a switch to flip – it's the tuner of your life, setting the station you live on. And there are two doors into that house:

- Bottom-Up Regulation – the door through the body.*
- Top-Down Regulation – the door through the mind.*

Bottom-Up Regulation begins in the body.

It's the language of movement, breath, touch, and sensation – primal whispers from muscle, fascia, and organs traveling upward to the brain. Each time you shake your arms, tap your sternum, or sink into a slow belly breath, you are stepping through the first door, telling your nervous system: You are safe. Science calls this afferent input – 80% of the vagus nerve's fibers carry these signals up to the brain, shaping mood, thoughts, and perception before you even think the words.

Top-Down Regulation begins in the mind.

It's the art of thought, focus, and meaning-making – the cortex choosing which story to tell the body. When you name your fear, speak a word like peace or enough, or reframe a racing thought, you are stepping through the second door, sending a signal down: Stand down. We are okay. This is efferent influence – the mind shaping heart rate, hormones, and the felt sense of safety.

Neither door is enough on its own.

Bottom-up softens the soil. Top-down plants the seed.

One clears the static. The other sets the station.

And so the invitation is simple: Remember.

Remember there are two doors – and you can walk through both.

Remember to weave the body and mind together.

Remember that safety isn't just thought – it's felt.

When you breathe as you choose new words, when you move as you shift old meanings, the system re-tunes. Fight, flight, and freeze fade. And the quiet, steady station comes through – the one you were always meant to live on: calm, connected, creative.

This reflection is the bridge into protocol integration. Every tool from here – breathwork, eye movements, somatic patterns, word anchors – will work best when you hold this truth close:

Healing has two doors. The magic happens when you open them both.

Part 9

Module 3

Clinical Integration and Case Studies

Module 3: Clinical Integration and Case Studies

Turning Knowledge Into Wisdom – The Art of Nervous System-Based Therapy

Module Overview

In Module 3, we move from understanding and technique into true clinical application. This module is about real clients, complex cases, and the nuanced decisions therapists must make when applying nervous system regulation in practice.

You've learned the science. You've practiced the techniques. Now, we explore:

- How to build full protocols for unique client presentations
- How to adapt and layer interventions over time
- What to do when clients don't respond as expected
- How to refine your clinical intuition using nervous system markers

Learning Objectives

By the end of this module, participants will be able to:

- Construct regulation-based intervention plans tailored to diagnosis and dysregulation patterns
- Identify when an intervention is mismatched to a client's state and adjust accordingly
- Apply nervous system-informed reasoning to troubleshoot challenging cases
- Reflect on their own nervous system role in the therapeutic relationship
- Confidently begin designing layered, diagnosis-specific protocols using breath, eye movement, somatic, and sensory strategies

Opening Reflection

“The nervous system is not a checklist. It’s a conversation.”

In real life, clients don’t arrive in neat categories. They arrive with burnout, pain, trauma, chaos, or collapse. Your job isn’t just to apply a technique — it’s to read the body’s language, match the state, and adjust the rhythm until something inside them says: Yes. Here. I can rest. I can reconnect.

This module is for clinicians who want to move beyond tools — into embodied therapeutic mastery.

Clinical Flowchart: How to Build a Nervous System-Based Protocol

Why a Flow Matters

Many therapeutic models focus on symptoms or diagnoses first. Nervous system regulation shifts the focus to state first, strategy second.

This flowchart offers a structured, state-based sequence that helps you build personalized, regulation-first protocols using the tools from Modules 1 and 2.

How to introduce it into your practice

Start with the first R – Recognize. Note all the ANS cues a client present with. Get comfortable with it and then move on to the next R – Respond and then R – Restore.

Nervous system work is a flow – get comfortable with it – do not force it.

Introduce one regulatory approach at a time if you feel more comfortable. Example: start with breathing, add breathing to your consultations as “warm-up”. Grow your protocols from there.

Add it as warm-up to every session or do it as a complete session.

Nervous System Protocol Flowchart

Step 1: Assess the Client's Current State

Use:

- Window of Tolerance Plotting Tool
- HRV symptoms checklist
- Verbal and non-verbal cues (e.g., breath, eyes, posture, reactivity)

Ask:

- Is the client in hyperarousal (panic, racing thoughts)?
- In hypoarousal (numb, foggy, shut down)?
- Or in functional freeze (masked, over-regulated)?

Step 2: Start with Stabilization Tools

Goal: Safety and Presence

Use:

- Grounding (feet, wall push, chair drop)
- Proprioceptive input
- Tactile anchoring
- Co-regulation through tone, rhythm, and pacing

Skip cognitive tasks until regulation baseline is present.

Step 3: Select Primary Regulation Entry

Match to state and symptom profile.

State	Entry Tools
Hyperarousal	Breathwork (Box, Coherence), Lateral Eye Tracking, Tactile Anchors
Hypoarousal	Upward Gaze, Vestibular Input, Cross-body Movement
Functional Freeze	Containment + Bilateral Tapping, Gentle Breath, Rhythmic Sway

Step 4: Observe Response and Adjust

Monitor:

- Breath pace, facial tone, orientation
- Verbal changes: more grounded, coherent speech
- Emotional shifts: tears, clarity, relief

If client dysregulates:

- Downshift (e.g., from breath to tactile)
- Return to Step 2 or reduce stimulation

Step 5: Layer Additional Tools Once stable

- Add bilateral input, walking, or symbolic work
- Consider interoception, reflective prompts, or memory integration
- For children: embed in play, rhythm, routine

Step 6: Track and Adjust Over Time

Weekly or monthly:

1. Reassess Window of Tolerance
2. Use HRV tracking or symptom logs
3. Update protocol flow based on state capacity and resilience markers

"Start with what the body says. Not what the story says."

Case Example 1: Generalized Anxiety & Functional Freeze

"I'm holding it together, but inside I'm drowning."

Client Profile

- Client: Female, 36 years old
- Occupation: Executive assistant in a corporate firm
- Diagnosis: Generalized Anxiety Disorder (GAD), perfectionism traits
- Presenting Symptoms:
 - Constant mental overdrive, difficulty sleeping
 - Frequent muscle tension, jaw clenching
 - Emotionally flat but tightly wound
 - Appears calm in session but admits to "meltdowns in private"

Nervous System State

- Functional Freeze (sympathetic overdrive masked by dorsal compensation)
- Low HRV coherence score on emWave (incoherence, despite parasympathetic markers)

BOLT score: 14 seconds (shallow, stress-based breathing)

Outcome After 3 Sessions

- Sleep improved from 4 to 6 hours/night
- Reduced jaw clenching; started journaling emotions
- Reports: "I don't just feel better. I feel like I'm in my body again."

Protocol Flow

Step	Tool or Technique	Notes
1. Stabilize & Orient	Chair anchoring + deep proprioceptive foot press	Gave client sense of boundary and “legs under her”
2. Entry Point: Tactile + Breath	Hand over chest + extended exhale (4:6)	Eyes stayed open; verbalized “I don’t feel safe closing my eyes”
3. Visual Regulation	Lateral eye tracking with soft object	Marked shift: shoulders softened, speech slowed
4. Emotional Emergence	Bilateral tapping with “It’s safe to feel now” mantra	Client cried for first time in 6 months of therapy
5. Integration & Homework	Daily breath tracking, object anchor in pocket	Began using tool in real-time work stress moments

Case Example 2: PTSD and Emotional Flooding

"It comes out of nowhere. I go from fine to drowning in a second."

Client Profile

- Client: Male, 28 years old
- Occupation: Firefighter, recently returned from medical leave
- Diagnosis: PTSD (work-related trauma), sleep disruption, hypervigilance
- Presenting Symptoms:
 - Sudden emotional outbursts, especially anger and tears
 - Nightmares, restlessness, intrusive memories
 - Easily startled, avoids eye contact
 - Feels "out of control" and ashamed of dysregulation

Nervous System State

- Clear hyperarousal dominance with sympathetic flooding
- Low coherence (HRV pattern erratic under breath), high resting HR
- BOLT score: 9 seconds — over-breathing, fast shallow inhale/exhale
- Unable to sit still for long without fidgeting

Outcome After 4 Sessions

- Reports fewer intrusive memories
- Sleep improved with regulation routine at bedtime

Protocol Flow

Step	Tool or Technique	Notes
1. Grounding & Containment	Weighted object on thighs + firm proprioceptive input (wall press)	Helped slow body activation and stabilize start of session
2. Breath + Visual Focus	Extended exhale (4:6) with eyes on fixed object	Client fidgeted initially, but settled by round 3
3. Emotional Discharge	Cross-body march + bilateral tapping	Tears + spontaneous memory retrieval; therapist held space
4. Coherence Training	2-minute coherence breath (HeartMath pacing)	Introduced after safety was established and emotions processed
5. Homework Tools	Sensory kit (cold object, texture, tapping), 5-min morning breath	Used daily after waking from nightmares or before shifts

Case Example 3: Child with Sensory Processing & Emotional Meltdowns

"Everything feels too much. Then I explode."

Client Profile

- Client: Boy, 7 years old
- Diagnosis: Sensory Processing Disorder (SPD), suspected autism spectrum traits
- Presenting Symptoms:
 - Daily meltdowns triggered by transitions, noise, or clothing textures
 - Avoids eye contact, sensitive to light and sound
 - Rigid routine needs; difficulty verbalizing feelings
 - Frequently toe-walks, struggles with body awareness

Nervous System State

- Highly sensory defensive, living in chronic sympathetic activation
- Alternates between hyperarousal and crash-like withdrawal
- Window of Tolerance is extremely narrow
- BOLT score (measured via child-friendly method): <6 seconds
- Avoids typical breathwork; responds better to movement-based regulation

Outcome After 5 Sessions

- Fewer meltdowns (from 5/week to 2/week)
- Began initiating rhythmic play during overstimulation
- Developed emotional "menu" visuals for identifying needs
- Parent reports: "We're learning to listen to his body, not just his words."

Protocol Flow

Step	Tool or Technique	Notes
1. Safe Space Setup	Tent corner with dim lighting + weighted items	Created predictable, low-stim environment before any intervention
2. Proprioceptive Input	Animal walks, wall push, and weighted blanket hugs	Introduced rhythm through body movement, not instructions
3. Rhythmic Eye Movement Play	"Spy Game" – tracking toy left to right, no verbal cue	Helped build tolerance for focus and co-regulation without pressure
4. Bilateral Regulation	Drum sticks: tap left/right with rhythm + music	Visible calming effect; added to home routine before school
5. Parent Integration	Co-regulation breathing while seated behind child (hand on back)	Taught parent to mirror nervous system calm using rhythm, not words

Case Example 4: Female Client with Burnout & Dysautonomia

"I'm tired all the time — but wired. Nothing helps me switch off."

Client Profile

- Client: Female, 44 years old
- Occupation: High-achieving business owner, mother of three
- Diagnosis: Burnout, suspected dysautonomia (POTS pattern), chronic fatigue
- Presenting Symptoms:
 - Constant exhaustion despite sleep
 - Elevated resting heart rate, dizziness upon standing
 - Irritability, mental fog, GI issues
 - Difficulty with deep breathing and structured meditation

Nervous System State

- Parasympathetic overactivity without coherence — likely dorsal vagal dominance
- HRV shows low variability with erratic baseline, low coherence
- BOLT score: 12 seconds — breath-holding causes dizziness
- Overstimulated by typical breathwork, responds paradoxically to extended exhales

Outcome After 4 Sessions

- Client reported: "I no longer feel like I'm failing at self-care"
- Energy stabilized mid-day; reduced dizzy spells
- Began using visual and somatic cues throughout her workday
- HRV coherence showed slight improvement; more important: client felt empowered, not exhausted by her tools

Protocol Flow

Step	Tool or Technique	Notes
1. Somatic Awareness First	Gentle swaying in seated position + tracking body tension	Helped shift from cognitive mode into present-moment body state
2. Proprioceptive Anchoring	Grounded walking barefoot + wall push during transitions	Introduced physical containment without breath manipulation
3. Visual + Sensory Input	Soft gaze tracking + cool cloth on back of neck	Stimulated vagus gently without overwhelming the system
4. Modified Breath Cueing	Gentle nasal breathing with no count, guided by rhythm not ratio	Removed pressure of performance; no breath holds used
5. Integration Tool	Created morning/evening self-regulation "menu" including movement, touch, visual orientation	Replaced strict routines with choice-based micropractices

Clinical Troubleshooting: When Techniques Fail or Backfire

"The right tool at the wrong time is still the wrong tool."

Why Some Regulation Tools Don't Work (Yet)

Even science-based techniques can backfire if the client is:

- In the wrong state for the tool
- Pushing beyond their current window of tolerance
- Lacking adequate co-regulation or safety
- Using the tool from a place of pressure or performance

Remember: Regulation is relational and sequential. Before skill comes readiness.

Clinical Mantras to Guide You

- "Start lower than you think, and slower than you want."
- "Coherence isn't comfort — it's capacity."
- "If it's not landing, it's not wrong — it's just not now."

Common Clinical Challenges & What To Do

Problem	What It May Mean	What To Try Instead
Client becomes dizzy with breathwork	Overbreathing, poor CO ₂ tolerance, or autonomic fragility	Use proprioception or visual tracking instead. Try sway + tactile input before returning to breath
Eye tracking triggers anxiety	Hypervigilance, unprocessed trauma, or visual-vestibular mismatch	Begin with grounding + peripheral vision games or softer visual engagement
Client "doesn't feel anything"	Dissociation, dorsal dominance, or low interoceptive awareness	Use textured objects, bilateral tapping, or somatic mirroring with therapist
Tool works once, then stops	Window of tolerance shifted, nervous system adapted	Reassess state; consider layering or switching entry system (e.g., from breath to movement)
Client reports shame about not "getting it right"	Performance anxiety, fawn response, or high cognitive override	Normalize setbacks, simplify instructions, return to co-regulation or playful tone

Final Integration: From Technique to Transformation

"It's not about adding more tools. It's about knowing when to use what you already have."

Integration Is Not Protocol—It's Presence

By now, you've learned:

- How to identify autonomic states using clinical markers
- How to apply breath, eye, somatic, and sensory strategies
- How to build individualized nervous system protocols
- How to troubleshoot and adapt with clinical intuition

But the deeper work is this: becoming a regulating presence yourself. A grounded, attuned therapist becomes the external nervous system until the client can self-regulate.

Therapist as Regulator

What makes the biggest difference in client outcomes?

Not just the tool.

Not just the plan.

But the therapist's ability to:

- Co-regulate in real time
- Sense when to pause, pivot, or deepen
- Hold presence instead of pushing for progress

Putting It All Together

Every client = a different entry point

Every session = a new state

Every technique = a doorway, not a destination

The art of this work is not in applying all the tools.

It's in listening deeply to the body's invitation — and responding wisely.

Suggested Daily Integration (for therapists)

Practice	Purpose
Begin your day with 3 mins of coherence breathing	Anchor yourself before entering session space
Track 1 client each week more deeply: breath, eyes, posture	Train your observational regulation lens
Reflect weekly: "What did I regulate well? What felt forced?"	Build clinical intuition through practice

Part 10

Appendixes

CPD Assessment:

Instructions

- This assessment aligns with HPCSA CPD requirements
- Please complete all questions clearly and independently
- A score of 70% or higher is required to qualify for certification
- Submit your completed answers to the course facilitator www.neurosyncacademy.com – use any of the email addresses provided on the website

SECTION A: Multiple Choice (1 mark each)

1. When building a nervous system regulation protocol, which should be assessed first?

- A. Cognitive function
- B. Symptom diagnosis
- C. Current autonomic state
- D. Preferred breathing style

2. What does a low BOLT score typically suggest?

- A. Elevated coherence
- B. Strong parasympathetic tone
- C. Poor CO₂ tolerance and over-breathing
- D. Increased vagal resilience

3. Functional freeze is best described as:

- A. Complete emotional shutdown
- B. Fight-or-flight dominance
- C. Blended sympathetic and dorsal vagal suppression
- D. Regulated hypoarousal

4. If a client becomes dizzy during extended exhale breathing, the therapist should:

- A. Encourage deeper breathing
- B. Add breath holds
- C. Switch to proprioceptive or visual tools
- D. End the session

5. Coherence training is most effective when:

- A. The client is in a high-alert sympathetic state
- B. The nervous system has been stabilized first
- C. Combined with hyperventilation techniques
- D. Used only during sleep routines

SECTION B: Short Answer

6. Describe how you would assess a client's readiness for breathwork using nervous system markers.

7. Give an example of a mismatch between a client's nervous system state and a technique — and how you would adjust your approach.

8. List 3 observable signs that suggest a regulation technique is working in real time.

9. Why is therapist regulation as important as technique selection in nervous system-based work?

10. Write a 4-step protocol for a client presenting with chronic anxiety, shallow breathing, and racing thoughts.

11. What are the three main branches of the autonomic nervous system?
12. What is coherence and why is it clinically relevant?
13. Name two signs of sympathetic nervous system dominance.
14. What is the ideal BOLT score range indicating good CO₂ tolerance?
15. Describe one breath technique that supports parasympathetic activation.

Section 2: Practical Demonstration Checklist

Instructions: illustrate with images

1. Coherence Breathing

2. Bilateral Eye Movement (horizontal tracking)

3. Grounding Practice (e.g., wall press, foot stomp)

4. Somatic Release (e.g., tapping or swaying)

5. Client State Identification (matching technique to nervous system state)

Section 3: Final Case-Based

1. A client reports dizziness with breathwork. What might this indicate, and what would you do instead?
2. A child becomes agitated during visual tracking. What regulation tools could you use instead?
3. List 3 clinical signs that suggest a client is moving into their window of tolerance.
4. Design a 3-step regulation protocol for a client with chronic anxiety.
5. Why is therapist regulation critical when applying nervous system tools?

BUSINESS & APPLICATION ADD-ON: How to Integrate This Model Into Your Practice

Translating into Your Clinical Work

This work is not designed as 'theory for theory's sake.

It is fully intended to be embedded into diverse clinical environments — whether you work individually, in groups, in hospitals, schools, or coaching environments.

Below are some ways to bring this into your professional work.

Clinical Application Models:

Individual Treatment Application

- Trauma-focused therapy
- Anxiety disorders
- Chronic pain rehabilitation
- PTSD stabilization
- Burnout recovery
- Executive coaching for stress regulation
- Occupational therapy physical engagement protocols
- Pediatric sensory regulation programs
- Vagal tone biofeedback integration

Group Therapy Applications

- Group stabilization track for inpatient psychiatric hospitals
- Anxiety groups with psychoeducation + regulation practice
- Burnout & resilience groups for medical professionals
- Trauma stabilization groups before trauma-processing groups
- Parent education groups on nervous system regulation for children
- Co-regulation groups for couples or family therapy

Hospital or Programmatic Use

- Incorporate into:
 - Inpatient trauma units
 - Pain management programs
 - Physical rehabilitation groups
 - Eating disorder stabilization protocols
 - DBT stabilization adjunct
 - Occupational therapy group models

Clinical Business Tip Box:

Business Model Tip:

You can create:

- Short 4–6 week stabilization intensives
- Ongoing nervous system maintenance groups
- Multi-disciplinary packages with psych, OT, physio, dietician partnerships

